

Analyzing variability and decomposing electricity-generation emission factors for three U.S. states

Zahra Ghaemi^a, Amanda D. Smith^{a,b},

^a*Department of Mechanical Engineering, University of Utah, Salt Lake City, UT 84112, USA*

^b*Pacific Northwest National Laboratory, Richland, WA 99352, USA*

Abstract

Electricity generation emission factors (EGEF) quantify the relationship between amount of pollutant emitted and amount of electricity generated. Quantifying variability among calculated EGEFs is important when EGEFs are used for decision-making. Variabilities in EGEF on an annual basis due to variability in the amount of coal, natural gas, and petroleum emissions within the fuel mix are quantified for California, Texas, and New York in 2017. The results show a higher coefficient of variation for SO_2 and NO_x compared to CO_2 EGEF. Changes in the EGEF over time are studied using decomposition analysis for California, Texas, and New York from 1990 to 2017. The results show that the main factor in reducing EGEF in California is the improving generation efficiency of power plants; in Texas, it is the increasing the ratio of renewable to non-renewable electricity generation; and in New York, it is the changing mix of fossil fuels that are consumed for electricity generation. The effect of variability of EGEF on environmental impact categories is analyzed. Eutrophication of air, eutrophication of water, and smog formation are subject to high uncertainty because SO_2 and NO_x EGEFs are used to quantify these impacts, whereas global warming potential has less uncertainty because it only uses CO_2 EGEF.

Keywords: electricity generation emission factor, fossil fuel emission factor, decomposition analysis, Monte Carlo simulation, environmental impact assessment

Nomenclature

AP	Acidification Potential
C	Emissions from electricity generation (kg)
C_{fuel}	Emission by each fossil fuel type (coal, natural gas, or petroleum) (kg)
EF	Emission factor
EGEF	Electricity generation emission factor
Eff	Efficiency
EIA	U.S. Energy Information Administration
EPA	U.S. Environmental Protection Agency
EA	Eutrophication of air
EW	Eutrophication of water
e_{fuel}	Emission factor for each fossil fuel (kg-emissions/MMBtu)
F_{fuel}	Energy input by each fossil fuel type (coal, natural gas, or petroleum) (MMBtu)
FF	Fossil fuel
G	Total electricity generation by state (MWh)

Email address: amanda.d.smith@utah.edu (Amanda D. Smith)

16	GWP	Global Warming Potential
17	HH	Human health
18	IDA	Index decomposition analysis
19	LMDI	Logarithmic mean divisia index analysis
20	m_{fuel}	The electricity generation from each fossil fuel divided by the total electricity generation
21		from all fossil fuels (coal, natural gas, and petroleum)
22	NG	Natural gas
23	Pet	Petroleum
24	PV	Photovoltaics
25	SA	Smog Air
26	U.S.	United States
27	u_{fuel}	The energy input divided by the electricity generation for each fossil fuel (MMBtu/MWh)
28	Q	Electricity generation by all fossil fuels (coal, natural gas, and petroleum) (MWh)
29	Q_{fuel}	Electricity generation by each fossil fuel type (coal, natural gas, or petroleum) (MWh)

30 1. Introduction

31 Climate change has ramifications on ecosystems, human health, and the energy sector ([National Oceanic](#)
32 [and Atmospheric Administration, 2019](#)). Human activities and energy consumption, on both global and
33 regional scales, are driving climate change ([Stocker et al., 2013](#)). The electricity sector alone contributed to
34 27.5% of 2017 greenhouse gas emissions in the United States (U.S.), making the sector the country’s second-
35 largest source of greenhouse gas emissions ([U.S. Environmental Protection Agency \(EPA\)](#)) and an essential
36 factor in reducing or mitigating climate change. Electricity generation influences ecosystems, human health,
37 and the climate by releasing CO_2 , SO_2 , and NO_x emissions through fossil fuel-fired generation. Monitoring
38 and measuring the greenhouse gas emissions resulting from electricity generation empowers a country or region
39 to move toward a low-carbon future ([Khan, 2019a](#)). However, the quantification of greenhouse gas emissions
40 resulting from electricity generation has been associated with a variety of uncertainties and variabilities that
41 may affect such decision-making and carbon management ([Li and Huang, 2012](#)). Therefore, defining and
42 quantifying these variabilities and changes in electricity generation-related emissions is also essential. This
43 study aims to define and quantify these factors accordingly.

44 In this work, we investigate variations between calculated electricity generation emission factor (EGEF)
45 values across U.S. states. Moreover, we quantify the key factors underlying changes in these EGEFs from
46 1990 to 2017. This work was conducted at the state level, and we have focused on three U.S. states to
47 illustrate a variety of cases. We chose California, Texas, and New York for this analysis because they are
48 three of the country’s most populous states. Also, they encompass a wide range of climate conditions and
49 sociopolitical systems, and more electricity sector-related data is publicly available for these three states
50 than for many other U.S. states. Moreover, each of these three states is located in one of the country’s
51 three major electrical grid interconnections. California lies within the Western Interconnection; the vast
52 majority of Texas lies within the Electric Reliability Council of Texas (ERCOT); and New York lies within
53 the Eastern Interconnection ([U.S. Department of Energy Office of Electricity \(2019\)](#)). These three states
54 have implemented progressive strategies to increase their renewable electricity generation and reduce their
55 electricity generation carbon footprints ([Barbose, 2017](#)).

56 This work addresses the following key research questions:

- 57 1. How do EGEFs (that is, the range of expected values for CO_2 , SO_2 , and NO_x emissions per unit of
58 electricity) vary across in three U.S. states of California, Texas, and New York?
- 59 2. How did these EGEFs change from 1990 to 2017, and what underlying factors drove these changes in
60 each of the studied states?

3. Can we reasonably quantify electricity generation’s environmental impacts at the state level, given the uncertainty surrounding the data used to calculate these states’ emission factors?

In the following subsections, we discuss the research background related to each of these key questions.

1.1. How do EGEFs (that is, the range of expected values for CO_2 , SO_2 , and NO_x emissions per unit of electricity) vary across in three U.S. states of California, Texas, and New York?

We define the term “emission factor” (EF) according to its use by the U.S. EPA: “An EF is a representative value that attempts to relate the quantity of a pollutant released to the atmosphere with an activity associated with the release of that pollutant” (U.S. EPA). A fossil fuel EF is the quantity of emissions per unit of energy content in a fuel (kg emissions/MMBtu), and this quantity is often provided as a single value; for example, the “EFs for greenhouse gas inventories” document by the EPA provides nationwide averages for coal, natural gas, and petroleum, which are responsible for 66%, 32%, and 1% of total U.S. CO_2 emissions for electricity generation (U.S. Energy Information Administration (EIA), 2019). In reality, a fossil fuel EF’s value varies year-to-year across different regions and types of combustion devices (Roy et al., 2009) due to the different efficiencies associated with power plants, various origins and sub-types of each fossil fuel used, and differences in the accuracy of emission measurements in each power plant.

EF value differences can significantly affect quantifications of CO_2 emissions (Liu et al., 2015). An electricity generation EF (EGEF) calculates the emissions released per unit of electricity generation (kg emissions/MWh generated). The greenhouse gas emissions associated with renewable and nuclear energy used to generate electricity are significantly lower and less variable than the emissions associated with electricity generation using fossil fuels: life-cycle greenhouse gas emissions have been reported at 45 (g CO_2 /kWh) for Photovoltaics (PV), 11 (g CO_2 /kWh) for wind, 12 (g CO_2 /kWh) for nuclear energy, and 1,000 (g CO_2 /kWh) for coal (National Renewable Energy Laboratory, 2013). Thus, PV, wind, and nuclear emissions’ impact on EGEF were neglected in the present study, and the EGEF can be quantified using a weighted sum of the EFs for the individual energy sources in a state’s fuel mix, multiplied by their consumption rates.

Several studies in the literature have investigated the effects of variabilities in the energy sources emissions on emissions accounting of the electricity generation.

Some studies analyzed variabilities in EGEF due to considering the marginal EGEF. Marginal EGEF is a metric to account for the changes in the electricity generation emissions as a result of an additional unit of electricity (Hawkes (2010) and Khan (2019a)). Studies that consider the marginal EGEF have shown variabilities in EGEF when the marginal EGEF is used compared to the average EGEF. For example, Siler-Evans et al. (2012) quantified the variabilities of marginal EGEF compared to the average EGEF for eight regional electricity generations. Their results showed that the marginal EGEFs were different from the average EGEFs up to 36% for the CO_2 EGEF, 280% for the SO_2 EGEF, and 161% for NO_x EGEF. Zheng et al. (2015) quantified the marginal EGEF and compared the marginal EGEF with the average EGEF for two case studies. Their results showed that the marginal EGEF was up to 50% higher than the average EGEF.

Some studies focused on the variabilities in the generation mix with seasons, days, and time of the day, and its effect on the EGEF (Khan, 2019b). Especially, the penetration of renewable sources and their intermittent electricity generation results in changes in the generation mix and variability of EGEF. For example, Khan et al. quantified the temporal variability of EGEF in Bangladesh (Khan, 2018) and New Zealand (Khan et al., 2018). They studied the variabilities in EGEF because of the changes in the electricity demand, the penetration of renewable sources, and the difference in the efficiency of power plants to generate electricity. Their results showed up to 40% variabilities in the EGEF. St-Jacques et al. (2020) studied the temporal EGEF of eight regions in the electrical zone of Ontario. They showed high variabilities in EGEF for different electrical zones compared to the average EGEF.

The mentioned studies perform an essential analysis in quantification and variability analysis of EGEF due to an additional unit of electricity demand and the changes in the time of electricity use. However, the EF of fossil fuels used to generate electricity can vary year-to-year due to the different efficiencies associated with power plants, various origins and sub-types of each fossil fuel used, and differences in the accuracy of emission measurements in each power plant. Hence, the annual variability of each fossil fuel EF changes the annual EGEF.

To the best of our knowledge, two studies performed variability analysis of fossil fuel EF to generate electricity.

Wang et al. (2019) used bootstrap and Monte Carlo simulations to determine methane EF variability for coal, using data from 11,000 power plants in China. They found high variability for methane EFs, which varied widely compared to the average EF values used in the 2006 Intergovernmental Panel on Climate Change (IPCC) “Guidelines for National Greenhouse Gas Inventories.” Roy et al. (2009) created equations for estimating coal EFs in India, finding this approach effective in reducing the uncertainty surrounding EFs for this fuel type. This approach yielded estimated EFs that were proposed to be more realistic than national-average EFs in representing coal EFs throughout India.

These studies have analyzed coal EF variability in emissions accounting in India and China. The literature has neglected, however, to analyze EGEFs’ variability, based on variabilities in each fossil fuel EFs at the U.S. state level, while considering the data’s limitations. Therefore, the current study’s first objective is to analyze EGEF’ variability with the available energy-source EFs at the state level, using our three selected states.

1.2. How did these emission factors change from 1990 to 2017, and what underlying factors drove these changes in each of the studied states?

EGEFs change over the years because of changes in the power generation process. In this work, we consider four major factors involved in those changes (Ang and Su (2016), Goh et al. (2018a), and Goh et al. (2018b)): (1) the fossil fuel share effect (i.e., fossil fuels’ share of electricity generation to non-fossil fuels—hydro-power, biomass, renewables, and nuclear—share of electricity generation), (2) the fossil fuel mix effect (i.e., the mix of fossil fuels consumed for electricity generation), (3) the generation efficiency effect (i.e., power plants’ generation efficiency), and (4) the EF effect (i.e., fossil fuel EFs). Understanding how EGEF can be broken down into these four factors can reveal the most effective factors in reducing EGEFs.

Decomposition analysis is a method of breaking the EGEF down into several predefined factors (Ang, 2004). Index decomposition analysis (IDA) is a useful method for performing decomposition analysis and evaluating the main forces driving changes in CO_2 emissions over the years (Xu and Ang (2013) and Zhang et al. (2013)).

Decomposition analysis of EGEFs has enabled a new understanding of the essential factors driving changes in EGEFs over the years. Ang and Su (2016) broke down the CO_2 EGEF in 10 Southeast Asian countries from 1990 to 2013. The results of this decomposition index identified the factor responsible for the largest reductions in several countries’ EGEFs: in Myanmar, this factor was an improvement in the country’s fossil fuel share effect; in Singapore and Thailand, this factor was changes in the countries’ fossil fuel mix effects; and in Vietnam, Brunei, and Malaysia, this factor was an improvement in power plants’ generation efficiency effect. Liu et al. (2017) used the IDA method to analyze the forces driving EGEFs at the provincial level in China. They concluded that an improvement to the generation efficiency effect was the most essential factor in reducing China’s EGEFs from 2000 to 2014.

Decomposition analysis of CO_2 emissions has also been conducted using electricity generation in the U.S. Jiang and Li (2017) used the LMDI (log mean Divisia index analysis) method to break down CO_2 emissions in the U.S. from 1990 to 2014, concluding that the country’s fossil fuel mix effect played the most significant role in reducing electricity CO_2 emissions from 1990 to 2005 (and in the CO_2 emissions predicted between 2005 and 2030). Mohlin et al. (2019) studied changes in CO_2 emissions at the U.S. state level from 1990 to 2015, showing that states’ fossil fuel share effect and fossil fuel mix effect were the main factors driving changes in CO_2 emissions.

The studies above demonstrated that the fossil fuel share effect, the fossil fuel mix effect, and the generation efficiency effect tend to account for the largest share of changes in EGEFs (Ang and Su (2016), Malla (2009), and Jiang and Li (2017)), and the most important factors differ between the national and provincial levels (Liu et al. (2017) and Mohlin et al. (2019)). Decomposition analysis of U.S. EGEFs is vital because magnitudes of greenhouse gas emissions resulting from electricity generation vary by location (de Chalendar et al., 2019) since U.S. states may belong to different electrical grid balancing authority areas, may have implemented different renewable energy policies (DSIRE), or may target different energy source mixes over the years while facing different public pressures to act upon these targets.

The current study builds on the work of Mohlin et al. (2019) by examining variability and different possible values for EGEFs, including changes in SO_2 and NO_x EGEF, extending the studied period to 2017, and focusing on three selected states that provide different case studies illustrating this variability. Thus,

this study’s second objective is to study EGEF changes over the years in California, Texas, and New York at the individual-state level.

1.3. Can we reasonably quantify electricity generation’s environmental impacts at the state level, given the uncertainty surrounding the data used to calculate these states’ emission factors?

If EGEFs are part of a decision-making process, then any metrics calculated using variations that affect EGEF values will also involve variability that is associated with the calculated metrics. The term “environmental impact,” as it is used here, describes the consequence of a plan, policy, or process. It is an example of a method used for decision- and policy-making. Environmental impact quantification is a method of evaluating a system or process’s environmental effects through four damage categories: human health, ecosystem quality, climate change, and resources (Jolliet et al., 2003). To quantify electricity generation’s environmental impacts, a computational tool can be used first to create a deterministic estimate. The variability associated with EGEF can be used with the analytical tool to quantify a resulting variability in environmental impacts. Traci, an open-source environmental impact assessment tool developed by the U.S. EPA (2012), was used in this study to quantify environmental impacts.

Many studies have quantified electricity generation’s environmental impact, but few analyses have incorporated uncertainty and variability into their results. Chen and Corson (2014) used a Monte Carlo simulation to determine the effects of EFs’ uncertainty in three environmental impact categories. They concluded that EFs’ variability and uncertainty could shift decision outcomes, based on environmental impact assessments. Sonnemann et al. (2003) estimated the effect of site-specific uncertainties and variability in measuring EGEF resulting from a waste incinerator, showing that significant uncertainties are associated with EGEFs because of inventory data uncertainties.

Tools that help decision-making can be improved if their results’ uncertainties and variabilities are more transparent (Tenney et al., 2006). We were unable to locate a study investigating the variability of a decision-making tool’s results, based on U.S. EGEFs’ variability. The lack of such a study may be due to a lack of reliable data on emissions’, EFs’, and EGEFs’ uncertainty and variability. Therefore, the current study’s third objective was to quantify how EGEFs’ variability affects the environmental impacts, using the limited data available from reliable sources.

Our analysis in this paper presents the following novel contributions:

1. The range and variability of EGEFs for CO_2 , SO_2 , and NO_x associated with fossil fuel electricity generation in the U.S. are quantified deterministically for all U.S. states. These EGEFs are quantified stochastically, using a Monte Carlo simulation for the three selected states.
2. The CO_2 EGEF is broken down for the three selected states, showing how those EGEFs have changed over time by revealing the main factors that have influenced changes to EGEFs between 1990 and 2017.
3. We quantify the variability of electricity generation’s environmental impacts due to variability in EGEFs. However, we caution against relying on any quantification of electricity generation’s environmental impacts at the U.S. state level.

2. Methodology

The following three subsections discuss the methods we used to address each of this study’s three objectives, respectively.

2.1. Emission factors’ distribution for electricity generation

This subsection explains our methodology in determining fossil fuel EFs’ variability in the U.S. in 2017 and then using that variability to identify the probability distribution of EGEFs at the state level in 2017. This subsection comprises three parts. First, the data sources and assumptions used to determine the EGEFs are described. Second, the formulations used to calculate EGEFs are presented. Third, the Monte Carlo simulation statistical tool is used to identify the results’ variability and probability distribution.

2.1.1. Data sources and assumptions

Three data sources from the “Electricity Detailed State Data from 1990 to 2017” webpage by the U.S. EIA were used to calculate energy-source EGEFs at the state level:

1. Annual data (from 1990 to 2018) for net generation by type of producer (e.g., the total electric power industry, electric generators, electric utilities, combined heat and power, and commercial power) and fuel type (e.g., coal, natural gas, petroleum, hydroelectric conventional, and wind) for 50 U.S. states.
2. Annual data (from 1990 to 2018) for fossil fuel consumption used for electricity generation by producer type (e.g., total electric power industry, electric generators, electric utilities, combined heat and power, and commercial power) and fuel type (e.g., coal, natural gas, petroleum, and other gases) for 50 U.S. states.
3. Annual data (from 1990 to 2017) for the electric power industry’s estimated emissions (CO_2 , SO_2 , and NO_x) for 50 U.S. states.

We made four assumptions in considering the fuels used for electricity generation at the state level:

1. Only emissions associated with fossil fuels’ combustion were considered, and emissions from the production, processing, transmission, and storage of energy sources were omitted from our calculations. We made this assumption because of the available electricity generation and emissions data from the EIA “Electricity Detailed State Data from 1990 to 2017” documents’ exclusion of other parts of the energy sources life cycle and focus on the production side of energy sources for electricity generation .
2. Emissions associated with renewables (hydro, wind, solar, biomass, and geothermal) and nuclear energy were also excluded from our calculations. Amponsah et al. (2014) reviewed greenhouse gas emissions at different stages of renewable technologies. In most cases (except for two instances using biomass technology), greenhouse gas emissions from power plant operations were less than 20 kg CO_{2eq} /MWh, which is less than greenhouse gas emissions associated with the use of fossil fuels to generate electricity (National Renewable Energy Laboratory, 2013).
3. “Other biomass” and “wood and wood derived fuels” energy sources from the “Electricity Detailed State Data from 1990 to 2017” documents were also excluded from our EGEF calculation. The three selected fossil fuels (i.e., coal, natural gas, and petroleum) are the top-three CO_2 emitters, accounting for 99% of CO_2 , 82% of SO_2 emissions, and 82% of NO_x emissions in U.S. electricity generation during 2017. The main energy source excluded from our analysis is “wood and wood derived fuels,” which account for 16% of SO_2 emissions; moreover, “other biomass” and “wood and wood derived fuels” respectively account for 8% and 5% of NO_x emissions. “Other biomass” and “wood and wood derived fuels” were omitted from this study because their consumption data is not available in the “Electricity Detailed State Data from 1990 to 2017” EIA documents.
4. The “other” fuel type was omitted because its heating value is unavailable, and it accounted for only 0.3% of total U.S. electricity generation in 2017, which is a marginal amount (“Electricity Detailed State Data from 1990 to 2017” EIA documents).

2.1.2. Electricity generation emission factor calculation

EF of fossil fuels can be calculated by dividing emissions at the state level for each fuel type by the amount of the consumption of that fuel to generate electricity (kg emissions/MMBtu), using Equation 1 (Intergovernmental Panel on Climate Change, 2006). EGEFs at the state level can be calculated using Equation 2, based on fuel consumption rates and fuel EFs, divided by electricity generation at the state level.

$$\text{Emission Factor}_{fuel} = \text{Emissions} / \text{Fuel Consumption} \quad (1)$$

$$\text{EGEF} = \frac{\sum_{fuel=1}^n \text{Fuel Consumption}_{fuel} \times \text{Emission Factor}_{fuel}}{\text{Electricity Generation}} \quad (2)$$

2.1.3. Statistical analysis

Probabilistic techniques, such as Monte Carlo simulation, can serve as a practical statistical tool in analyzing variability (Environmental Protection Agency, 1997). This study used the five-step Monte Carlo simulation procedure described by Hiraishi et al. (2001) to quantify EGEFs' variability at the state level.

1. Specify source-category uncertainties; fossil fuel EFs (kg CO_2 /MMBtu, kg SO_2 /MMBtu, and kg NO_x /MMBtu) for coal, natural gas, and petroleum are variable in the U.S.
2. Set up the necessary information: the quantities of CO_2 , SO_2 , and NO_x emissions, fossil fuel consumption rates, and electricity generation are available in the "Electricity Detailed State Data from 1990 to 2017" document by the EIA for all 50 U.S. states in 2017. Fossil fuel EFs' probability density functions were computed using the available continuous distributions in the SciPy Python library (SciPy library, 2019).
3. Select random variables: random CO_2 , SO_2 , and NO_x EF variables (kg CO_2 /MMBtu, SO_2 /MMBtu, or NO_x /MMBtu) were generated for coal, natural gas, and petroleum.
4. Estimate EGEFs: EGEFs were calculated for Texas, California, and New York using fossil fuel consumption rates (MMBtu) multiplied by fossil fuel EFs' random variable (kg CO_2 /MMBtu, SO_2 /MMBtu, or NO_x /MMBtu), using Equation 2.
5. Iterate and monitor the results: the EGEF value from Step 4 was stored, and then the process was repeated from Step 3.

These five steps were simulated using continuous distributions in the SciPy library in Python, and 10,000 samples from fossil fuel EFs were generated to show EGEFs' variability for the study's three emission types at the state level.

2.2. The most influential elements of electricity generation emission factors

This subsection describes the study's methodology to break EGEFs down into influential factors at the state level from 1990 to 2017.

Two main methods were used for this decomposition analysis: structural decomposition analysis (SDA), which uses input-output models, and IDA, which implements index number theory (Goh et al., 2018b).

IDA has been successfully applied in the literature to evaluate different factors' impacts on energy and changes in emissions over the years (Zhang et al., 2013). In 2004, after 25 years of research on decomposition analysis, researchers had not reached an agreement on the best decomposition method (Ang, 2004). Popular decomposition methods belong to two main groups: (1) methods linked to the Laspeyres index (i.e., the impact of a factor is calculated while all the other factors are constant at their base year) and (2) methods linked to the Divisia index (i.e., a weighted sum of logarithmic growth rates is used) (Ang et al., 1998).

The advantage of the Divisia index approach (i.e., the concept of logarithmic change) over the Laspeyres index approach (i.e., the familiar concept of percentage changes) is the Divisia index symmetric and additive features of relative changes (Ang, 2004). The symmetric feature of the Divisia index means that, in quantifying relative changes between two numbers, the order of the numbers does not affect relative change values. For example, an absolute value of $\log(1/5)$ equals $\log(5/1)$, but a percentage change between these two numbers gives values of 5 and 0.20 (1/5). The additive feature of the Divisia index means that two relative changes can be summed or subtracted. For example, $\log(5/1)$ equals $\log(5/3)$ added to $\log(3/1)$. But, using the concept of percentage changes, 5 does not equal 1.67 (5/3) added to 3, which gives 4.67.

For cross-country or regional analysis, conventional decomposition methods leave significant residuals, and perfect decomposition methods (e.g., LMDI)—which leave no residuals—have been suggested instead (Zhang and Ang, 2001). Ang (2004) concluded that LMDI is the most "elegant" decomposition method, from a theoretical perspective, for use in regional studies. Therefore, considering its additive and symmetric features and its leaving no residuals, we selected the LMDI method for the current study.

To apply the LMDI method to changes in EGEFs from this study's base year (1990) to its final year (2017) at the state level, Equation 3 was used as the governing equation to be broken down into four factors

299 (Ang and Su, 2016).

$$EGEF = \frac{C}{G} = \sum_{fuel=1}^3 \frac{Q}{G} \frac{Q_{fuel}}{Q} \frac{F_{fuel}}{Q_{fuel}} \frac{C_{fuel}}{F_{fuel}} \quad (3)$$

300	C	Emissions from electricity generation (kg)
301	G	Total electricity generation by state (MWh)
302	Q	Electricity generation by all fossil fuels (coal, natural gas, and petroleum) (MWh)
303	Q_{fuel}	Electricity generation by each fossil fuel type (coal, natural gas, or petroleum) (MWh)
304	F_{fuel}	Energy input by each fossil fuel type (coal, natural gas, or petroleum) (MMBtu)
305	C_{fuel}	Emission by each fossil fuel type (coal, natural gas, or petroleum) (kg)

306 Equation 3 can be rewritten as Equation 4 (Ang and Su, 2016).

$$EGEF = \sum_{fuel=1}^3 p \times m_{fuel} \times u_{fuel} \times e_{fuel} \quad (4)$$

307	p	The proportion of electricity produced from fossil fuels in the total electricity generation (MWh/MWh)
308	m_{fuel}	The electricity generation from each fossil fuel divided by the total electricity generation from
309		all fossil fuels (coal, natural gas, and petroleum) (MWh/MWh)
310	u_{fuel}	The energy input divided by the electricity generation for each fossil fuel generation (MMBtu/MWh)
311	e_{fuel}	The EGEF of each fossil fuel (kg-emissions/MMBtu)

312 Next, the total change in EGEFs at the state level from 1990 (the base year) to 2017 was shown as ΔV_{total}
 313 (kg emissions/MWh) and broken down into ΔV_p (fossil fuel share effect), ΔV_m (fossil fuel mix effect), ΔV_u
 314 (generation efficiency effect), and ΔV_e (EF effect) (kg emissions/MWh) factors in Equation 5.

$$\Delta V_{total} = V^T - V^0 = \Delta V_p + \Delta V_m + \Delta V_u + \Delta V_e \quad (5)$$

315 Further equations for the LMDI method are available in Appendix (also, see Ang et al. (1998), Ang
 316 (2005), Ang et al. (2011), and Ang and Su (2016) for detailed methodology).

317 2.3. The effects of electricity generation emission factors' present variability on environmental impact quan- 318 tification

319 This subsection describes the methodology we used to quantify the effects of EGEFs' variabilities on
 320 environmental impact categories during 2017 in California, Texas, and New York, using the five-step Monte
 321 Carlo procedure presented earlier under Subsection 2.1.

322 Environmental impact quantification anticipates a system or process's environmental effects through four
 323 damage categories: human health, ecosystem quality, climate change, and resources (Jolliet et al., 2003).
 324 Midpoint categories can be used to connect emission inventory data to damage categories. Six midpoint
 325 categories were used in the current study: global warming potential (GWP), acidification potential (AP),
 326 human health (HH), eutrophication air (EA), eutrophication water (EW), and smog air (SA). These midpoint
 327 categories were chosen because they connect CO_2 , SO_2 , and NO_X —the available emissions data in this
 328 study—to the damage categories. GWP affects climate change. AP, EA, EW, and SA affect ecosystem
 329 quality. Finally, HH and SA affect human health (Jolliet et al., 2003).

330 Traci, an open-source environmental impact assessment tool developed by the EPA, was used in this
 331 study to determine the environmental impacts of electricity generation at the state level by providing the
 332 characteristic factors shown in Table 1. Traci was selected for this study as a free, U.S.-based tool developed
 333 by the EPA, a reliable source for environmental studies.

334 To quantify the impact categories, first, the characteristic factors (Table 1) are collected for CO_2 , SO_2 ,
 335 and NO_X from the Traci tool; second, these three characteristic factors are multiplied by the three EGEF
 336 of CO_2 , SO_2 , and NO_X ; and third, the sum of the three multiplied values results in the equivalent emission

Characteristic factor	CO_2	SO_2	NO_x
GWP (kg CO_2 -eq/kg substance)	1	0	0
AP (kg SO_2 -eq/kg substance)	0	1	0.7
HH (kg $PM_{2.5}$ -eq/kg substance)	0	0.061	0.007
EA (kg N -eq/kg substance)	0	0	0.044
EW (kg N -eq/kg substance)	0	0	0.291
SA (kg O_3 -eq/kg substance)	0	0	24.79

Table 1: Characteristic factors of CO_2 , SO_2 , and NO_x for six impact categories (GWP, AP, HH, EA, EW, and SA) from the U.S. EPA (2012).

of each impact category. For example, to quantify the AP (SO_2 -eq) using the Traci tool, the second row of Table 1 is used. This row shows that the characteristic factor of CO_2 is zero (kg SO_2 -eq/kg substance), SO_2 is 1 (kg SO_2 -eq/kg substance), and NO_x is 0.7 (kg SO_2 -eq/kg substance) for AP. Hence, the CO_2 EGEF (lb CO_2 /MWh) is multiplied by zero, SO_2 EGEF (lb SO_2 /MWh) is multiplied by 1 (kg SO_2 -eq/kg substance), and NO_x EGEF (lb NO_x /MWh) is multiplied by 0.7 (kg SO_2 -eq/kg substance). Their sum results in each state’s AP of EGEF (lb SO_2 -eq/MWh).

A Monte Carlo simulation was used to calculate the variability of electricity generation’s environmental impacts, following the five-step procedure described in Subsection 2.1.

1. Specify source category uncertainties: EGEFs are variable (kg CO_2 /MWh, kg SO_2 /MWh, and kg NO_x /MWh) in California, Texas, and New York, based on the results of the first objective identified in Subsection 3.1.
2. Set up the necessary information: characteristic factors of CO_2 , SO_2 , and NO_x for six impact categories (GWP, AP, HH, EA, EW, and SA) were obtained from the Traci, which are shown in Table 1. EGEFs’ probability density functions can be calculated using the Subsection 2.1 method and are shown in Table 5 in Appendix.
3. Select random variables: random variables for CO_2 , SO_2 , and NO_x EGEF (kg CO_2 /MWh, SO_2 /MWh, or NO_x /MWh) were determined for California, Texas, and New York.
4. Estimate the environmental impact: GWP, AP, HH, EA, EW, and SA were calculated for California, Texas, and New York using the Traci characteristic factors in Table 1, multiplied by CO_2 , SO_2 , and NO_x EGEF random variables from Step 3.
5. Iterate and monitor the results: The environmental impact categories’ values from Step 4 were stored, and then the process was repeated from Step 3 to show the variability of EGEFs’ environmental impacts in California, Texas, and New York during 2017.

3. Results and Discussion

The results obtained by addressing each of this study’s three objectives are discussed in the three subsections below.

3.1. Emission factors’ distribution for electricity generation

This subsection first reveals the CO_2 , SO_2 , and NO_x EF distributions of coal, natural gas, and petroleum in the U.S. during 2017. Next, the variability of CO_2 , SO_2 , and NO_x EGEFs are presented for California, Texas, and New York in 2017.

The EF units for fossil fuels are compatible with the units in the “EFs for greenhouse gas inventories” document by the EPA, kg of emissions per MMBtu of fossil fuel consumption.

The probability distribution of CO_2 EFs for coal, natural gas, and petroleum are shown in Figures 1, 2, and 3, and their statistics are given in Table 2. The median of CO_2 EF using coal was 95.38 kg- CO_2 /MMBtu (0.32 kg- CO_2 /kWh), while the corresponding median for natural gas was 56.38 kg- CO_2 /MMBtu (0.19 kg- CO_2 /kWh), and the corresponding median for petroleum was 73.92 kg- CO_2 /MMBtu (0.25 kg- CO_2 /kWh). The average results for coal, natural gas, and petroleum EF were 95.52, 53.06, and 73 kg- CO_2 /MMBtu (0.33, 0.18, and 0.25 kg- CO_2 /kWh), respectively, in the “EFs for greenhouse gas inventories” document by

Type of EF	Distribution	Std dev (kg/mmBTU)	Mean (kg/mmBTU)	Median (kg/mmBTU)	Coefficient of variation %
CO_2 of Coal	Levy Stable	10.22	98.01	95.38	10.43
CO_2 of Natural gas	Power Log-Normal	2.63	56.71	56.38	4.65
CO_2 of Petroleum	Johnson SU	8.53	77.13	73.92	11.06
SO_2 of Coal	Folded Cauchy	0.06	0.09	0.08	61.56
SO_2 of Natural gas	Cauchy	0.00	0.00	0.00	36.25
SO_2 of Petroleum	Pearson Type III	0.17	0.16	0.11	109.08
NO_x of Coal	Folded Cauchy	0.03	0.07	0.07	38.12
NO_x of Natural gas	Folded Cauchy	0.10	0.07	0.03	155.71
NO_x of Petroleum	Johnson SB	0.07	0.11	0.10	66.42

Table 2: Statistics of CO_2 , SO_2 , and NO_x EFs of coal, natural gas, and petroleum in the U.S. during 2017

the EPA. A coefficient of variation (i.e., the data’s standard deviation, divided by the data’s mean) can be used to compare uncertainty across data sets when means differ. The coefficients of variation for CO_2 EF using coal, natural gas, and petroleum were 10.43%, 4.65%, and 11.06%, respectively.




Figure 1: Probability distribution of CO_2 EFs for coal in the 50 U.S. states during 2017.




Figure 2: Probability distribution of CO_2 EFs for natural gas in the 50 U.S. states during 2017.




Figure 3: Probability distribution of CO_2 EFs for petroleum in the 50 U.S. states during 2017.

The probability distribution of SO_2 EFs for coal, natural gas, and petroleum are shown in Figures A.1, A.2, and A.3 in Appendix, and their statistics are given in Table 2. The median of SO_2 EF using coal was 0.08 kg- SO_2 /MMBtu (0.27 g- SO_2 /kWh), while the corresponding median for natural gas was 0.0 kg- SO_2 /MMBtu (0.0 g- SO_2 /kWh), and the corresponding median for petroleum was 0.11 kg- SO_2 /MMBtu (0.37 g- SO_2 /kWh). The coefficient of variation for SO_2 EFs using coal, natural gas, and petroleum were 61.56%, 36.25%, and 109.08%, respectively.

The probability distribution of NO_x EFs for coal, natural gas, and petroleum are shown in Figures A.4, A.5, and A.6 in Appendix, and their statistics are given in Table 2. The median of NO_x EFs using coal was 0.07 kg- NO_x /MMBtu (0.24 g- NO_x /kWh), while the corresponding median for natural gas was 0.03 kg- NO_x /MMBtu (0.10 g- NO_x /kWh), and the corresponding median for petroleum was 0.10 kg- NO_x /MMBtu (0.34 g- NO_x /kWh). The coefficient of variation for NO_x EFs using coal, natural gas, and petroleum were 38.12%, 155.71%, and 66.42%, respectively.

The probability density functions of these nine EF types are depicted in Figures 1–3 and A.1–A.6, and




Figure 4: CO_2 EGEF distribution in Texas during 2017, using sources' (i.e., coal, natural gas, and petroleum) EF distributions

Type of EGEF	Distribution	Std dev (lb/kWh)	Mean (lb/kWh)	Median (lb/kWh)	Coefficient of variation %
CO_2 in California	Generalized Extreme	19.07	403.58	400.6	4.73
CO_2 in Texas	Tukey Lambda	97.03	1291.67	1291.25	7.51
CO_2 in New York	Generalized Extreme	17.77	391.96	389.22	4.53
SO_2 in California	Logarithm of the Gamma	0.05	0.01	0.0	767.39
SO_2 in Texas	Hyperbolic Secant	28.12	2.57	0.8	1093.1
SO_2 in New York	Pearson Type III	0.33	0.05	0.03	697.78
NO_x in California	Generalized Half-Logistic	6.81	0.65	0.25	1043.14
NO_x in Texas	Logarithm of the Gamma	16.25	2.13	0.91	764.04
NO_x in New York	Generalized Half-Logistic	6.33	0.63	0.25	1000.39

Table 3: Statistics of CO_2 , SO_2 , and NO_x EGEFs of California, Texas, and New York during 2017

their statistics are shown in Table 2. In Step 3 of this study's Monte Carlo method, random variables from probability density functions were generated and combined, based on each state's share of coal, natural gas, and petroleum (i.e., energy mix). In Step 4 of this study's Monte Carlo method, 10,000 samples were generated to calculate the CO_2 , SO_2 , and NO_x EGEFs. The probability distribution of EGEFs using CO_2 , SO_2 , and NO_x of the study's 10,000 samples are shown in Figures 4, A.7, and A.8 for Texas.

The CO_2 , SO_2 , and NO_x EGEFs in Texas were, respectively, 1,166 lb CO_2 /MWh, 1.3 lb SO_2 /MWh, and 0.8 lb NO_x /MWh in 2017, as per the "State Electricity Profiles in 2017" EIA document. The median values—using Figures 4, A.7, and A.8—were, respectively, 1,293.25 lb CO_2 /MWh, 0.83 lb SO_2 /MWh, and 0.9 lb NO_x /MWh, shown in Table 3, reflecting 10.9%, 36%, and 12.5% respective differences between the median values in this study and the data in the "Electricity Detailed State Data from 1990 to 2017" EIA document. The coefficient of variation (i.e., the data's standard deviation, divided by the data's mean) is shown in Table 3 for CO_2 , SO_2 , and NO_x EGEFs in Texas, which are 7%, 697%, and 1061%, respectively.

Figures A.9, A.10, and A.11 in Appendix show the distributions of CO_2 , SO_2 , and NO_x EGEFs in California.

The CO_2 , SO_2 , and NO_x EGEFs in California were, respectively, 474 lb CO_2 /MWh, 0.0 lb SO_2 /MWh, and 0.7 lb NO_x /MWh in 2017, as per the "State Electricity Profiles in 2017" EIA document. The median values—using Figures A.9, A.10, and A.11—were, respectively, 400.76 lb CO_2 /MWh, 0.0 lb SO_2 /MWh, and 0.24 lb NO_x /MWh, shown in Table 3, reflecting 15.4%, 0%, and 95% respective differences between the median values in this study and the data in the "Electricity Detailed State Data from 1990 to 2017" EIA document. The coefficient of variation is shown in Table 3 for CO_2 , SO_2 , and NO_x EGEFs in California, which are 4.8%, 505%, and 1444%, respectively.

Figures A.12, A.13, and A.14 in Appendix show the distribution of CO_2 , SO_2 , and NO_x EGEFs in

New York. The CO_2 , SO_2 , and NO_x EGEFs in New York were, respectively, 439 lb CO_2 /MWh, 0.3 lb SO_2 /MWh, and 0.5 lb NO_x /MWh in 2017, as per the “State Electricity Profiles in 2017” EIA document. The median values—using Figures A.12, A.13, and A.14—were, respectively, 389.22 lb CO_2 /MWh, 0.03 lb SO_2 /MWh, and 0.25 lb NO_x /MWh, shown in Table 3, reflecting 11.3%, 90%, and 50% differences between the median values in this study and the data in the “Electricity Detailed State Data from 1990 to 2017” EIA document. The coefficient of variation is shown in Table 3 for CO_2 , SO_2 , and NO_x EGEFs in New York, which are 4.6%, 699%, and 1,000%, respectively.

The variabilities in the EGEFs of this study are due to the variabilities in the EF of each energy source to generate electricity. Therefore, the main factor that causes the variabilities in EGEF of this study is different from the main factor that causes the variabilities in studies that quantified marginal EGEF or temporal EGEF in the literature. However, comparing the EGEF variabilities of studies in the literature with variabilities in this study can show the range and magnitude of variabilities of EGEF in the literature. Hawkes (2010) stated the marginal EGEF in the Great Britain of 1521 lb- CO_2 /MWh for 2002–2009, with error bars at 10%. Also, IPCC’s report on EGEF of power plants to generate electricity is associated with 1% to 10% uncertainty in CO_2 EGEFs. Therefore, the coefficient of variations of this study (4.73% for California, 7.51% for Texas, and 4.53% for New York) for CO_2 EGEF is in a similar range of uncertainties of variabilities in EGEFs in the literature.

Two conclusions can be drawn from a comparison of this study’s results for the three states examined:

1. For the three states, SO_2 and NO_x EGEFs showed significant variability compared to CO_2 EGEFs.
2. EGEFs for states with higher shares of coal, natural gas, and petroleum were estimated to differ less from the EIA document values. The shares of emissions in Texas, California, and New York were as follows:
 - (a) In Texas, 99% of CO_2 , 94% of SO_2 , and 94% of NO_x emissions for electricity generation resulted from coal, natural gas, and petroleum consumption, respectively, which resulted in the lowest difference between this study’s calculated values and the EIA document values.
 - (b) In California, coal, natural gas, and petroleum accounted for 95% of CO_2 emissions, 45% of SO_2 emissions, and 73% of NO_x emissions, respectively, for electricity generation. Including other energy sources (i.e., “wood and wood derived fuels,” “other biomass,” and “other”) would improve estimations of SO_2 and NO_x EGEFs in California.
 - (c) In New York, coal, natural gas, and petroleum were responsible for 97% of CO_2 emissions, 71% of SO_2 , and 63% of NO_x emissions for electricity generation, respectively. Including other energy sources (i.e., “wood and wood derived fuels,” “other biomass,” and “other”) would improve estimations of SO_2 and NO_x EGEFs in New York.

3.2. The most influential elements of electricity generation emission factors

This subsection breaks down four influential elements (the fossil fuel share effect, the fossil fuel mix effect, the generation efficiency effect, and the EF effect) of EGEFs to show how these four elements changed EGEFs at the state level from 1990 to 2017.

The LMDI method was performed for the study’s three examined states: Texas, California, and New York. The results of this process for the fossil fuel share effect, the fossil fuel mix effect, the generation efficiency effect, the EF effect, and the sum of all four effects (i.e., the total) are shown in Figure 5.

Figure 5 shows that total EGEFs have decreased in all three examined states from 1990 to 2017. The highest reduction occurred in New York, and the lowest reduction occurred in California.

The fossil fuel share effect was responsible for 42%, 36%, and 37% of the EGEF changes from 1990 to 2017 in Texas, California, and New York, respectively. This finding shows that fossil fuels’ share of electricity generation in the three states has decreased, substituted by renewables, nuclear energy, and biomass. From 1990 to 2017, coal, natural gas, and petroleum’s shares of electricity generation changed from 92% to 75% in Texas, from 50% to 43% in California, and from 61% to 38% in New York.

The fossil fuel mix effect was responsible for 15%, 36%, and 63% of the EGEF changes in Texas, California, and New York, respectively, from 1990 to 2017. This change in electricity EGEFs reveals that the three states’ fuel mixes changed to environmentally friendly fossil fuels by decreasing the shares of coal and petroleum and increasing the shares of natural gas from 1990 to 2017. Coal and petroleum’s shares of electricity generation




Figure 5: A comparison of Texas, California, and New York's respective changes in CO_2 EGEFs from 1990 to 2017 (lb CO_2)/MWh

reduced by 13% and 1%, respectively, in Texas, 2% and 3% in California, and 18% and 25% in New York. Natural gas shares increased by 2% in California and 20% in New York.

The generation efficiency effect was responsible for 45%, 60%, and 19% of the EGEF changes in Texas, California, and New York, respectively, from 1990 to 2017. This change means that more electricity was generated from one unit of fossil fuel over the years because of improved efficiency for power plants or generators. The definition of “generation efficiency” is compatible with the heat rate of fossil fuels (i.e., the energy input to a system, divided by electricity generation). The heat rate values of coal, natural gas, and petroleum have improved over the years in the U.S. (U.S. Energy Information Administration, 2019b). Changes to the generation efficiency effect with heat rate values are shown in Figures 1, 2, and 3 in the supplementary materials for California, Texas, and New York.

The EF effect increased in the three studied states, which shows that more emissions were produced from one unit of fossil fuel over the years. This increase in the EF effect might have resulted from these states' using more exact metrics.

The “Annual energy outlook 2019: with projections to 2040” EIA document described several elements that have influenced EGEFs. Three of these elements that can be quantified over the years were selected for observation in this study: natural gas prices, levelized costs of solar and wind energy, and old power plants' retirement capacity. Other elements can also change EGEFs but were not considered in this study. The selected three factors are part of a system, interconnecting with one another. The following connections show, to some extent, how these three elements might affect EGEFs over the years, though they share no simple cause-and-effect relationship.

Natural gas prices alter the fossil fuel mix effect by decreasing coal-fired power generation's competitiveness (the “Annual energy outlook 2019: with projections to 2040” EIA document). Therefore, it is expected to see lower EGEFs of fossil fuel mix effect with lower natural gas prices. Figures 4, 5, and 6 in the supplementary materials show the CO_2 EGEFs of the fossil fuel mix effect and natural gas prices in California, Texas, and New York from 1990 to 2017.

Levelized costs of electricity of solar and wind power decreased, based on a Lazard report, from 2009 to 2017. Such policies and renewables' LCOE helped EGEFs for the fossil fuel share effect decrease over the years at the state level. Figures 7, 8, and 9 in the supplementary materials show the renewable LCOE and fossil fuel share effects in California, Texas, and New York from 1990 to 2017.

The retirement of older and less-efficient power plants effectively reduced EGEFs at the state level over the years (U.S. EIA, 2019). Figures 10, 11, and 12 in the supplementary materials show the capacity for coal-, natural gas-, and petroleum-based power plants' retirement during each studied year in California, Texas, and New York, respectively.

Type	Unit	Std dev	Mean	Median	Coefficient of variation %
GWP of California	lb CO_2 -eq/MWh	19.07	403.58	400.6	4.73
GWP of Texas	lb CO_2 -eq/MWh	97.03	1291.67	1291.25	7.51
GWP of New York	lb CO_2 -eq/MWh	17.77	391.96	389.22	4.53
AP of California	lb SO_2 -eq/MWh	4.77	0.46	0.18	1028.66
AP of Texas	lb SO_2 -eq/MWh	30.31	4.06	1.58	746.23
AP of New York	lb SO_2 -eq/MWh	4.44	0.49	0.21	905.02
HH of California	lb $PM_{2.5}$ -eq/MWh	0.05	0.01	0.0	961.4
HH of Texas	lb $PM_{2.5}$ -eq/MWh	1.72	0.17	0.06	997.87
HH of New York	lb $PM_{2.5}$ -eq/MWh	0.05	0.01	0.0	665.49
EA of California	lb N -eq/MWh	6.81	0.65	0.25	1043.14
EA of Texas	lb N -eq/MWh	16.25	2.13	0.91	764.04
EA of New York	lb N -eq/MWh	6.33	0.63	0.25	1000.39
EW of California	lb N -eq/MWh	1.98	0.19	0.07	1043.14
EW of Texas	lb N -eq/MWh	4.73	0.62	0.26	764.04
EW of New York	lb N -eq/MWh	1.84	0.18	0.07	1000.39
SA of California	lb O_3 -eq/MWh	168.77	16.18	6.12	1043.14
SA of Texas	lb O_3 -eq/MWh	402.88	52.73	22.55	764.04
SA of New York	lb O_3 -eq/MWh	156.83	15.68	6.19	1000.39

Table 4: Statistics of GWP, AP, HH, EA, EW, and SA of California, Texas, and New York during 2017

3.3. The effects of electricity generation emission factors' present variability on environmental impact quantification

This subsection discusses the variabilities of six environmental impact categories—global warming potential (GWP), acidification potential (AP), human health (HH), eutrophication air (EA), eutrophication water (EW), and smog air (SA)—based on variability in EGEFs in California, Texas, and New York in 2017.

The results' SO_2 and NO_x characteristics for GWP, which are shown in Table 1, stood at zero. Thus, CO_2 EGEFs drove GWP in this study. The median value of Texas's GWP (1292 [lb CO_2 -eq/MWh]) exceeded California's GWP (401 [lb CO_2 -eq/MWh]) and New York's GWP (389 [lb CO_2 -eq/MWh]), which are shown in Table 4. This finding is compatible with the higher CO_2 EGEFs of Texas compared to California and New York, which are shown in Table 3. As we have shown, a coefficient of variation (i.e., the data's standard deviation divided by the data's mean) can be used to compare data sets' uncertainty when means differ. This study's coefficients of variation for GWP were 4.8%, 7.4%, and 4.5 % for California, Texas, and New York, respectively. These values equal the coefficient of variation for CO_2 EGEFs in Table 3 since they represent the same data's coefficient of variation.

Acidification potential was calculated based on SO_2 emissions (with a characteristic factor of 1 [kg SO_2 -eq/kg substance]) and NO_x emissions (with a characteristic factor of 0.7 [kg SO_2 -eq/kg substance]). The median value of Texas's AP (1.58 [lb SO_2 -eq/MWh]) exceeded California's AP (0.18 [lb SO_2 -eq/MWh]) and New York's AP (0.21 [lb SO_2 -eq/MWh]), which are shown in Table 4. This finding is compatible with the higher SO_2 and NO_x EGEFs of Texas compared to California and New York shown in Table 3. The coefficient of variation for AP were 1029%, 746%, and 905% for California, Texas, and New York, respectively.

Human health's impact was calculated based on SO_2 emissions (with a characteristic factor of 0.061 [$PM_{2.5}$ -eq/kg substance]) and NO_x emissions (with a characteristic factor of 0.007 [$PM_{2.5}$ -eq/kg substance]). The median value of Texas's HH (0.06 [lb SO_2 -eq/MWh]) exceeded California's HH (0.0 [lb [$PM_{2.5}$ -eq/MWh]) and New York's AP (0.0 [lb [$PM_{2.5}$ -eq/MWh]), which are shown in Table 4. This finding is compatible with the higher SO_2 and NO_x EGEFs of Texas compared to California and New York shown in Table 3. The coefficient of variation for HH were 767%, 1093%, and 697% for California, Texas, and New York, respectively.

The eutrophication of air, water, and smog formation were calculated based on NO_x EGEFs, with characteristic factors of 0.044 kg N-eq/kg substance for the eutrophication of air, 0.291 kg N-eq/kg substance for the eutrophication of water, and 24.793 kg O_3 -eq/kg substance for the smog formation. The coefficients of variabilities for each state were constant across the three categories because of the constant variability of NO_x EGEF for each state, which equaled 1043%, 764%, and 1000% (shown in Table 4) for California, Texas, and New York, respectively. These values equal the coefficient of variation for NO_x EGEFs in Table 3 since they represent the same data's coefficient of variation.

GWP, which represents CO_2 EGEFs' variability, had the lowest coefficient variability among the assessed environmental categories. The coefficients of variability for EA, EW, and SA were constant, as NO_x EGEFs' variability remained the same for each state. Also, EA, EW, and SA had the highest coefficient variability

among the environmental categories examined. These results show that higher variabilities are associated with environmental impacts that use SO_2 and NO_x EGEFs.

Figures A.17– A.22 shows the statistical visualization of showing the changes in the minimum, maximum, median, and range of the six environmental impact categories (i.e., GWP, AP, HH, EA, EW, and SA) in California, Texas, and New York from 1990 to 2017.

4. Limitations and future work

This work involved the following limitations:

- We assumed the three fossil fuel types—coal, natural gas, and petroleum—were used identically in different states. This assumption was incorrect; different sub-types of coal, natural gas, and petroleum are used for electricity generation. Also, fossil fuels’ characteristics differ, depending on the fuels’ region of extraction, as data for the 50 U.S. states, categorized into these three main fossil fuel types, in the EIA’s “Electricity Detailed State Data from 1990 to 2017” document reveal.
- Moreover, we neglected “wood and wood derived fuels,” “other biomass,” and “other” in quantifying EGEFs in this study since their consumption rate data were unavailable in the “Electricity Detailed State Data from 1990 to 2017” EIA document. However, “Electricity Detailed State Data from 1990 to 2017” EIA document on electricity emissions stated “wood and wood derived fuels” is responsible for 16% of SO_2 emissions and 5% NO_x emissions from electricity generation in the U.S. Also, “other biomass” is responsible for 8% of NO_x emissions from electricity generation in the country.
- The characteristic factors of quantifying environmental impacts, shown in Table 1, are also site-specific, depending on a region of study. In this study, an average number for the three U.S. states that we considered was obtained since data on site-specific characteristics were unavailable in the U.S., to the best of our knowledge.
- CO_2 emission, electricity generation, and energy source consumption data were available for approximately 6,000 power plants throughout the U.S., which enabled a calculation of CO_2 EGEFs. EGEFs are more concentrated and show less variability when the data derives from power plants, instead of states. However, a limitation of using data from power plants is that SO_x and NO_x emissions are unavailable in the data sets within the “Electricity emissions by plant and by region” EIA document, and only CO_2 EGEFs could be calculated. This problem limited our estimation of SO_x and NO_x EGEFs, as well as our calculations of the variability in environmental impacts from different categories involving SO_x and NO_x EGEFs.

The potential topics for future work following up on the current study are as follows.

- A consumption based method of accounting electricity generation emissions should be considered in future work since the current study focused on the production side. A consumption based accounting system considers the emissions from outside the system boundary (Ghaemi and Smith, 2020) and considers how changes in demand affect EGEFs. It also includes upstream (i.e., fossil fuel extraction, refinery, material manufacturing, and construction) and downstream (i.e., marketing, transmission, and distribution) emissions from electricity generation .
- In the current study, variability in environmental impacts and electricity generation were limited to variability in EGEFs from fossil fuels. Measurement uncertainties regarding fossil fuel consumption, emissions, and electricity generation reveal more information on variability in environmental impacts and electricity generation.
- Four factors were broken down from the current study’s governing equation of electricity generation, using the LMDI method. Other factors could be broken down from the governing equation using the LMDI method, such as sector-specific factors, to reveal more information about how EGEFs have changed in California, Texas, and New York.

5. Conclusions

As we have shown, the year-to-year variability in fossil fuel emission factors (EF) in the 50 U.S. states has caused variations in the annual quantities of electricity generation emission factors (EGEF) at the state level. Knowing the range of expected values for EGEFs enables an understanding of the decision risks associated with fossil fuel EFs. The relative ranges (i.e., the ratios of the ranges and the means of the data set) for CO_2 EGEFs were 43%, 59%, and 42% in California, Texas, and New York, respectively, during 2017. The coefficients of variation or relative standard deviations (i.e., the ratio of standard deviation and the mean of the data set) were 4.7%, 7.4%, and 4.5% for CO_2 EGEFs in California, Texas, and New York, respectively, in 2017. These variabilities in EGEFs have shown that using site-specific fossil fuel EFs is necessary when making important decisions using EGEFs.

Moreover, this study by using the background data from Energy Information Administration (EIA) has shown that EGEFs changed from 1990 to 2017 in the U.S. Four elements that affect these EGEF changes can be broken down from the governing equation (Equation 3) for electricity generation: (1) the fossil fuel share effect (i.e., the share of fossil fuel to non-fossil fuel [hydro-power, biomass, renewables, and nuclear energy] in electricity generation), (2) the fossil fuel mix effect (i.e., the mix of fossil fuels consumed for electricity generation), (3) the generation efficiency effect (i.e., power plants' generation efficiency), and (4) the EF effect (i.e., the EFs of energy sources). California, Texas, and New York reduced their EGEFs from 1990 to 2017 by 100, 227, and 316 $lbCO_2/MWh$, respectively. The most effective element of this reduction in New York was the fossil fuel mix effect, which reduced the state's EGEFs by 198 $lbCO_2/MWh$. In California, the generation efficiency effect decreased the state's EGEFs by 60 $lbCO_2/MWh$. In Texas, the EGEF has been mitigated by 130 $lbCO_2/MWh$ because of changes in the fossil fuel share effect. These results indicate that state-specific laws and regulations should be enacted to effectively mitigate EGEFs since each state's potential to reduce electricity generations emissions differs.

Variability and uncertainty in EGEFs affect the decision-making tools' results. One such decision-making tool used in the emissions reduction field is environmental impact assessments. EGEFs are used to calculate electricity generation's environmental impact, which shows the variability in environmental impact results. In this study, global warming potential showed the lowest variability among such results, with relative standard deviations of 4.8%, 7.4%, and 4.5% in California, Texas, and New York, respectively, in 2017. Eutrophication of air, water, and smog formation (based on NO_x emissions) showed the highest variability because of EGEFs' variability. herefore, calculated environmental impacts will vary significantly when considering SO_2 or NO_x EGEFs compared with considering CO_2 EGEFs. As a result, decision-makings that use eutrophication of air, water, and smog formation categories are unreasonable due to dealing with relatively high uncertainty compared to decision-makings that use global warming potential.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

CRedit authorship contribution statement

Zahra Ghaemi: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Software, Visualization, Writing - original draft, and Writing - review & editing. **Amanda D. Smith:** Conceptualization, Methodology, Project administration, Supervision, and Writing - review & editing.

Acknowledgements

We appreciate the assistance and support provided by the University of Utah Department of Facilities Management.

Data Availability

Three data sources from the “Electricity Detailed State Data from 1990 to 2017” webpage by the U.S. Energy Information Administration were used to calculate electricity generation emission factors at the U.S. state level from 1990 to 2017 in this paper:

1. The annual data for net electricity generation (Net Generation by State by Type of Producer by Energy Source (EIA-906, EIA-920, and EIA-923) excel file)
2. The annual data for fossil fuel consumption used for electricity generation (Fossil Fuel Consumption for Electricity Generation by Year, Industry Type and State (EIA-906, EIA-920, and EIA-923) excel file)
3. The annual data for the electric power industry’s estimated emissions (CO_2 , SO_2 , and NO_x) (U.S. Electric Power Industry Estimated Emissions by State (EIA-767, EIA-906, EIA-920, and EIA-923) excel file)

References

- N. Y. Amponsah, M. Trolborg, B. Kington, I. Aalders, R. L. Hough, Greenhouse gas emissions from renewable energy sources: A review of lifecycle considerations, *Renewable and Sustainable Energy Reviews* 39 (2014) 461–475.
- B. Ang, B. Su, Carbon emission intensity in electricity production: A global analysis, *Energy Policy* 94 (2016) 56–63.
- B. Ang, P. Zhou, L. Tay, Potential for reducing global carbon emissions from electricity production—a benchmarking analysis, *Energy Policy* 39 (5) (2011) 2482–2489.
- B. W. Ang, Decomposition analysis for policymaking in energy:: which is the preferred method?, *Energy policy* 32 (9) (2004) 1131–1139.
- B. W. Ang, The LMDI approach to decomposition analysis: a practical guide, *Energy policy* 33 (7) (2005) 867–871.
- B. W. Ang, F. Zhang, K.-H. Choi, Factorizing changes in energy and environmental indicators through decomposition, *Energy* 23 (6) (1998) 489–495.
- G. Barbose, US renewables portfolio standards: 2017 annual status report, Tech. Rep., Lawrence Berkeley National Lab.(LBNL), Berkeley, CA (United States), 2017.
- X. Chen, M. S. Corson, Influence of emission-factor uncertainty and farm-characteristic variability in LCA estimates of environmental impacts of French dairy farms, *Journal of cleaner production* 81 (2014) 150–157.
- J. A. de Chalendar, J. Taggart, S. M. Benson, Tracking emissions in the US electricity system, *Proceedings of the National Academy of Sciences* 116 (51) (2019) 25497–25502.
- Environmental Protection Agency, Policy For Use Of Probabilistic Analysis In Risk Assessment, 1997.
- Z. Ghaemi, A. D. Smith, A review on the quantification of life cycle greenhouse gas emissions at urban scale, *Journal of Cleaner Production* 252 (2020) 119634.
- T. Goh, B. Ang, B. Su, H. Wang, Drivers of stagnating global carbon intensity of electricity and the way forward, *Energy Policy* 113 (2018a) 149–156.
- T. Goh, B. Ang, X. Xu, Quantifying drivers of CO_2 emissions from electricity generation—Current practices and future extensions, *Applied Energy* 231 (2018b) 1191–1204.
- A. D. Hawkes, Estimating marginal CO_2 emissions rates for national electricity systems, *Energy Policy* 38 (10) (2010) 5977–5987.

661 T. Hiraishi, B. Nyenzi, J. Penman, S. Habetsion, Quantifying uncertainties in practice, IPCC Good Practice
662 Guidance and Uncertainty Management in National Greenhouse Gas Inventories .

663 Intergovernmental Panel on Climate Change, 2006 IPCC guidelines for national greenhouse gas inventories,
664 Intergovernmental Panel on Climate Change, 2006.

665 X.-T. Jiang, R. Li, Decoupling and decomposition analysis of carbon emissions from electric output in the
666 United States, Sustainability 9 (6) (2017) 886.

667 O. Jolliet, M. Margni, R. Charles, S. Humbert, J. Payet, G. Rebitzer, R. Rosenbaum, IMPACT 2002+: a new
668 life cycle impact assessment methodology, The international journal of life cycle assessment 8 (6) (2003)
669 324.

670 I. Khan, Importance of GHG emissions assessment in the electricity grid expansion towards a low-carbon
671 future: A time-varying carbon intensity approach, Journal of Cleaner Production 196 (2018) 1587–1599.

672 I. Khan, Greenhouse gas emission accounting approaches in electricity generation systems: A review, Atmo-
673 spheric Environment 200 (2019a) 131–141.

674 I. Khan, Temporal carbon intensity analysis: renewable versus fossil fuel dominated electricity systems,
675 Energy Sources, Part A: Recovery, Utilization, and Environmental Effects 41 (3) (2019b) 309–323.

676 I. Khan, M. W. Jack, J. Stephenson, Analysis of greenhouse gas emissions in electricity systems using time-
677 varying carbon intensity, Journal of Cleaner Production 184 (2018) 1091–1101.

678 Y. Li, G. Huang, Electric-power systems planning and greenhouse-gas emission management under uncer-
679 tainty, Energy conversion and management 57 (2012) 173–182.

680 N. Liu, Z. Ma, J. Kang, A regional analysis of carbon intensities of electricity generation in China, Energy
681 Economics 67 (2017) 268–277.

682 Z. Liu, D. Guan, W. Wei, S. J. Davis, P. Ciais, J. Bai, S. Peng, Q. Zhang, K. Hubacek, G. Marland, et al.,
683 Reduced carbon emission estimates from fossil fuel combustion and cement production in China, Nature
684 524 (7565) (2015) 335–338.

685 S. Malla, CO_2 emissions from electricity generation in seven Asia-Pacific and North American countries: a
686 decomposition analysis, Energy Policy 37 (1) (2009) 1–9.

687 K. Mohlin, A. Bi, S. Brooks, J. Camuzeaux, T. Stoerk, Turning the corner on US power sector CO_2 emis-
688 sions—a 1990–2015 state level analysis, Environmental Research Letters 14 (8) (2019) 084049.

689 National Oceanic and Atmospheric Administration, Climate change impacts, [https://www.noaa.gov/
690 education/resource-collections/climate-education-resources/climate-change-impacts](https://www.noaa.gov/education/resource-collections/climate-education-resources/climate-change-impacts), ac-
691 cessed: 2020-01-07, 2019.

692 National Renewable Energy Laboratory, Life Cycle Greenhouse Gas Emissions from Electricity Generation .

693 J. Roy, P. Sarkar, S. Biswas, A. Choudhury, Predictive equations for CO_2 emission factors for coal combustion,
694 their applicability in a thermal power plant and subsequent assessment of uncertainty in CO_2 estimation,
695 Fuel 88 (5) (2009) 792–798.

696 SciPy library, Continuous distributions in SciPy library in Python, [https://docs.scipy.org/doc/scipy/
697 reference/stats.html](https://docs.scipy.org/doc/scipy/reference/stats.html), accessed: 2021-02-04, 2019.

698 K. Siler-Evans, I. L. Azevedo, M. G. Morgan, Marginal emissions factors for the US electricity system,
699 Environmental science & technology 46 (9) (2012) 4742–4748.

700 G. W. Sonnemann, M. Schuhmacher, F. Castells, Uncertainty assessment by a Monte Carlo simulation in
701 a life cycle inventory of electricity produced by a waste incinerator, Journal of cleaner production 11 (3)
702 (2003) 279–292.

- M. St-Jacques, S. Bucking, W. O'Brien, Spatially and temporally sensitive consumption-based emission factors from mixed-use electrical grids for building electrical use, *Energy and Buildings* 224 (2020) 110249.
- T. F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, P. M. Midgley, et al., *Climate change 2013: The physical science basis*, 2013.
- A. Tenney, J. Kværner, K. I. Gjerstad, Uncertainty in environmental impact assessment predictions: the need for better communication and more transparency, *Impact Assessment and Project Appraisal* 24 (1) (2006) 45–56.
- U.S. Department of Energy Office of Electricity, Learn more about the interconnections, <https://www.energy.gov/oe/services/electricity-policy-coordination-and-implementation/transmission-planning/recovery-act-0>, accessed: 2020-03-26, 2019.
- U.S. EIA, Annual energy outlook 2019: with projections to 2040, <https://www.eia.gov/outlooks/ieo/pdf/ieo2019.pdf>, accessed: 2019-12-10, 2019.
- U.S. Energy Information Administration, Average Tested Heat Rates by Prime Mover and Energy Source, 2008 - 2018, https://www.eia.gov/electricity/annual/html/epa_08_02.html, accessed: 2020-02-03, 2019b.
- U.S. Energy Information Administration, Electricity emissions by plant and by region, <https://www.eia.gov/electricity/data/emissions/>, accessed: 2019-12-09, 2019c.
- U.S. Energy Information Administration, State Electricity Profiles in 2017, <https://www.eia.gov/electricity/state/archive/2017/>, accessed: 2020-02-03, 2019a.
- U.S. Energy Information Administration (EIA), Electricity Detailed State Data from 1990 to 2017, <https://www.eia.gov/electricity/data/state/>, accessed: 2019-09-09, 2019.
- U.S. Environmental Protection Agency, Emission factors for greenhouse gas inventories, https://www.epa.gov/sites/production/files/2018-03/documents/emission-factors_mar_2018_0.pdf, accessed: 2019-10-23, 2018.
- U.S. Environmental Protection Agency, Basic Information of Air Emissions Factors and Quantification, <https://www.epa.gov/air-emissions-factors-and-quantification/basic-information-air-emissions-factors-and-quantification#About%20Emissions%20Factors>, accessed: 2020-01-27, 2019b.
- U.S. Environmental Protection Agency, inventory of U.S. greenhouse gas emissions and sinks 1990-2017, <https://www.epa.gov/sites/production/files/2019-04/documents/us-ghg-inventory-2019-main-text.pdf>, accessed: 2020-01-07, 2019a.
- U.S. EPA, Tool for Reduction and Assessment of Chemicals and Other Environmental Impacts (TRACI), https://www.epa.gov/sites/production/files/2015-12/traci_2_1_2014_dec_10_0.xlsx, accessed: 2019-11-26, 2012.
- K. Wang, J. Zhang, B. Cai, S. Yu, Emission factors of fugitive methane from underground coal mines in China: Estimation and uncertainty, *Applied Energy* 250 (2019) 273–282.
- X. Xu, B. W. Ang, Index decomposition analysis applied to CO_2 emission studies, *Ecological Economics* 93 (2013) 313–329.
- F. Zhang, B. Ang, Methodological issues in cross-country/region decomposition of energy and environment indicators, *Energy Economics* 23 (2) (2001) 179–190.
- M. Zhang, X. Liu, W. Wang, M. Zhou, Decomposition analysis of CO_2 emissions from electricity generation in China, *Energy policy* 52 (2013) 159–165.
- Z. Zheng, F. Han, F. Li, J. Zhu, Assessment of marginal emissions factor in power systems under ramp-rate constraints, *CSEE Journal of Power and Energy Systems* 1 (4) (2015) 37–49.




Figure A.1: Probability distribution of SO_2 EFs for coal in the 50 U.S. states during 2017.




Figure A.2: Probability distribution of SO_2 EFs for natural gas in the 50 U.S. states during 2017.




Figure A.3: Probability distribution of SO_2 EFs for petroleum in the 50 U.S. states during 2017.




Figure A.4: Probability distribution of NO_x EFs for coal in the 50 U.S. states during 2017.




Figure A.5: Probability distribution of NO_x EFs for natural gas in the 50 U.S. states during 2017.

ΔV_{total} shows the total impact of changes in EGEFs in each state from 1990 (the base year) to 2017.
 ΔV_p , ΔV_m , ΔV_u , and ΔV_e also relate to the impact of changes in p , m_{fuel} , u_{fuel} , e_{fuel} .

$$\Delta V_{total} = V^T - V^0 = \Delta V_p + \Delta V_m + \Delta V_u + \Delta V_e \quad (A.1)$$

Following the implementation of the LMDI method from [Ang and Su \(2016\)](#), ΔV_p , ΔV_m , ΔV_u , and ΔV_e can be calculated using Equations A.2, A.3, A.4, and A.5, respectively.

$$\Delta V_p = \sum_{fuel=1}^3 \frac{w^T_{fuel} - w^0_{fuel}}{\ln \frac{w^T_{fuel}}{w^0_{fuel}}} \ln \frac{p_j^T}{p_j^0} \quad (A.2)$$




Figure A.6: Probability distribution of NO_x EFs for petroleum in the 50 U.S. states during 2017.

$$\Delta V_m = \sum_{fuel=1}^3 \frac{w^T_{fuel} - w^0_{state,fuel}}{\ln \frac{w^T_{fuel}}{w^0_{fuel}}} \ln \frac{m_j^T}{m_j^0} \quad (A.3)$$

$$\Delta V_u = \sum_{fuel=1}^3 \frac{w^T_{fuel} - w^0_{fuel}}{\ln \frac{w^T_{fuel}}{w^0_{fuel}}} \ln \frac{u_j^T}{u_j^0} \quad (A.4)$$

$$\Delta V_e = \sum_{fuel=1}^3 \frac{w^T_{fuel} - w^0_{fuel}}{\ln \frac{w^T_{fuel}}{w^0_{fuel}}} \ln \frac{e_j^T}{e_j^0} \quad (A.5)$$

752 where w^T_{fuel} and w^0_{fuel} in Equation A.6 were used.

$$w^T_{fuel} = \frac{C^T_{fuel}}{G^T} \quad w^0_{fuel} = \frac{C^0_{fuel}}{G^0} \quad (A.6)$$

Type	CO_2	SO_2	NO_x
Coal EF in the U.S.	Levy Stable	Folded Cauchy	Folded Cauchy
Natural Gas EF in the U.S.	Power Log-Normal	Cauchy	Folded Cauchy
Petroleum EF in the U.S.	Johnson SU	Pearson Type III	Johnson SB
EGEF in California	Generalized Extreme	Logarithm of the Gamma	Generalized Half-Logistic
EGEF in Texas	Tukey Lambda	Hyperbolic Secant	Logarithm of the Gamma
EGEF in New York	Generalized Extreme	Pearson Type III	Generalized Half-Logistic

Table A.1: EGEF and fossil fuel EFs' distribution types for CO_2 , SO_2 , and NO_x




Figure A.7: SO_2 EGEF distribution in Texas during 2017, using sources' (i.e., coal, natural gas, and petroleum) EF distributions




Figure A.8: NO_x EGEF distribution in Texas during 2017, using sources' (i.e., coal, natural gas, and petroleum) EF distributions




Figure A.9: *CO₂* EGEF distribution in California during 2017, using sources' (i.e., coal, natural gas, and petroleum) EF distributions




Figure A.10: *SO₂* EGEF distribution in California during 2017, using sources' (i.e., coal, natural gas, and petroleum) EF distributions




Figure A.11: NO_x EGEF distribution in California during 2017, using sources' (i.e., coal, natural gas, and petroleum) EF distributions




Figure A.12: CO_2 EGEF distribution in New York during 2017, using sources' (i.e., coal, natural gas, and petroleum) EF distributions




Figure A.13: SO_2 EGEF distribution in New York during 2017, using sources' (i.e., coal, natural gas, and petroleum) EF distributions




Figure A.14: NO_x EGEF distribution in New York during 2017, using sources' (i.e., coal, natural gas, and petroleum) EF distributions




Figure A.15: A comparison of Texas, California, and New York's changes in SO_2 EGEFs from 1990 to 2017 (10^{-3} -lb SO_2)/MWh.




Figure A.16: A comparison of Texas, California, and New York's changes in NO_x EGEFs from 1990 to 2017 (10^{-3} -lb NO_x)/MWh.




Figure A.17: A statistical visualization of GWP (lb CO_2 -eq/MWh) showing the minimum, maximum, and median of the California, Texas, and New York from 1990 to 2017




Figure A.18: A statistical visualization of AP (lb SO_2 -eq/MWh) showing the minimum, maximum, and median of the California, Texas, and New York from 1990 to 2017




Figure A.19: A statistical visualization of HH (lb $PM_{2.5}$ -eq/MWh) showing the minimum, maximum, and median of the California, Texas, and New York from 1990 to 2017




Figure A.20: A statistical visualization of EA (lb N-eq/MWh) showing the minimum, maximum, and median of the California, Texas, and New York from 1990 to 2017




Figure A.21: A statistical visualization of EW (lb N -eq/MWh) showing the minimum, maximum, and median of the California, Texas, and New York from 1990 to 2017




Figure A.22: A statistical visualization of SA (lb O_3 -eq/MWh) showing the minimum, maximum, and median of the California, Texas, and New York from 1990 to 2017