

Partial Molar Volume of NaCl and CsCl in Mixtures of Water and Methanol by Experiment and Molecular Simulation

Maximilian Kohns¹, Martin Horsch, Hans Hasse

*Laboratory of Engineering Thermodynamics (LTD), University of Kaiserslautern,
Erwin-Schrödinger Str. 44, D-67663 Kaiserslautern, Germany*

Abstract

Densities of solutions of NaCl and CsCl in mixtures of water and methanol are determined by experiment and molecular dynamics simulation. Both experiments and simulations cover the concentration range up to the solubility limit of the salt in the temperature range $288.15 \leq T / \text{K} \leq 318.15$ at ambient pressure. Non-polarizable molecular models from the literature are used for the ions and solvents. The partial molar volume of the salts at infinite dilution in the mixed solvent is determined from an empirical correlation of the data. The mixed solvent effects on the density and the partial molar volumes of the salts are well predicted by the molecular models.

Keywords: density measurement, molecular simulation, mixed solvent,

¹Author to whom correspondence should be addressed. Electronic mail: maximilian.kohns@mv.uni-kl.de. Telephone: +49-631/205-3216. Fax: +49-631/205-3835.

1 **1. Introduction**

2 A sound knowledge of thermophysical properties of electrolyte solutions is
3 important for the understanding of natural processes and the design of a
4 wide variety of industrial processes. Aqueous electrolyte solutions have been
5 studied extensively, and more recently, also non-aqueous electrolyte solutions
6 have drawn significant attention, because of their importance e.g. in energy
7 storage. However, solutions of salts in mixed solvents have been studied only
8 rarely up to now. In the present work, we therefore investigate the den-
9 sity and related volumetric properties in mixed solvent electrolyte solutions,
10 where the electrolytes are either NaCl or CsCl and the solvent is a mixture
11 of water and methanol.

12 Several experimental studies on densities of solutions of alkali halide salts in
13 mixtures of water and methanol have been reported in the literature. Take-
14 naka et al. [1–3] report densities of such solutions for seven of the 17 alkali
15 halide salts that are soluble both in water and in methanol, namely LiCl,
16 NaCl, KCl, NaBr, KBr, NaI, and KI. For solutions of NaCl, there are three
17 additional investigations [4–6], and Ivanov and Abrosimov [7] study solutions
18 of KBr. Furthermore, there is a detailed investigation by Raatschen [8] on

19 solutions of LiBr in mixtures of water and methanol. Werblan [5] reports
20 data on the cesium halides, however, inspection of these data shows consid-
21 erable scatter, cf. Section 4.3.

22 Interestingly, densities of solutions of electrolytes in mixtures of water and
23 methanol have hardly been studied with thermodynamic models. There are
24 several modeling studies dealing with phase equilibria in such systems [9–
25 19], however, most of these works employ models for the excess Gibbs
26 energy, so that solution densities cannot be obtained. Equations of state
27 (EOS), which enable density calculations, have only very recently been ex-
28 tended towards the modeling of mixed solvent electrolyte solutions. This
29 is e.g. the case for the ePC-SAFT EOS [17], the SAFT-VRE EOS [18] and
30 the electrolyte CPA EOS [19]. Molecular simulations are particularly attrac-
31 tive for modeling mixed solvent electrolyte solutions due to the low number
32 of adjustable parameters and a strong physical background. However, we
33 are aware only of the work of Strauch and Cummings [12], who investigate
34 the vapor-liquid equilibrium of solutions of NaCl in mixtures of water and
35 methanol and also report the densities of the coexisting phases.

36 In contrast, solutions of a salt - in most cases NaCl - in pure water have been
37 studied extensively with molecular simulations in the recent literature. In

38 these studies, mainly the differences between polarizable and non-polarizable
39 models were investigated. Models including polarizability, such as the alkali
40 halide models developed around the polarizable water models SWM4-DP
41 [20] and BK3 [21], were found to perform better than non-polarizable ones in
42 some aspects, e.g. concerning the description of activity coefficients [22, 23].
43 Interestingly, this does not necessarily result in an improved prediction of the
44 salt solubility [23]. When considering other properties such as the density,
45 simple non-polarizable models are found to be of almost the same quality
46 as the polarizable ones [23]. Furthermore, non-polarizable models can in
47 principle describe a variety of properties of electrolyte solutions fairly well.
48 However, the systematic assessment of Orozco et al. [24] shows that none of
49 the existing model parameterizations is able to quantitatively describe sev-
50 eral properties at once. This suggests that better model parameterizations
51 can be found [25]. For a more detailed discussion of the recent advances in
52 the field, see the comprehensive review by Nezbeda et al. [26].

53 In the present work, we first report new experimental density data for the
54 systems water-methanol-NaCl and water-methanol-CsCl. Our experiments
55 cover the entire concentration range up to the solubility limit of the salt,
56 and we report data for the temperatures 288.15, 293.15, 298.15, 308.15, and

57 318.15 K. Second, we address the question whether simple non-polarizable
58 molecular models based on Lennard-Jones (LJ) sites and partial charges can
59 predict volumetric properties of the studied electrolyte solutions. In our dis-
60 cussion, the partial molar volume of the salt at infinite dilution is of particu-
61 lar interest, because it is a very sensitive property and provides a descriptive
62 view of the salt-solvent interactions.

63 Throughout this work, the composition of a ternary solution of a salt CA,
64 which completely dissociates into the ions C^+ and A^- , in a mixture of wa-
65 ter and methanol is described by the methanol mole fraction of the salt-free
66 solvent mixture χ_{MeOH}

$$\chi_{\text{MeOH}} = n_{\text{MeOH}} / (n_{\text{MeOH}} + n_{\text{W}}) \quad (1)$$

67 and the true mole fraction of the cation x_{C^+}

$$x_{C^+} = n_{C^+} / (n_{C^+} + n_{A^-} + n_{\text{MeOH}} + n_{\text{W}}), \quad (2)$$

68 where n_i are the mole numbers.

69 2. Experiments

70 Ultradry methanol (≤ 50 ppm water) with a purity of ≥ 99.9 % was pur-
71 chased from Roth. Deionized water was produced by an Elix Essential 5^{UV}
72 of Merck Millipore and degassed by boiling before use. NaCl was purchased
73 from Merck with a purity of ≥ 99.5 % and CsCl was purchased from Roth
74 with a purity of ≥ 99.999 %. The salts were dried in a vacuum oven at 353 K
75 for 24 h.

76 The electrolyte solutions were prepared gravimetrically (AE240, Mettler-
77 Toledo) in a glovebox (GS Glovebox Technik). In case of mixed solvents,
78 about 200 ml of a stock solution of water + methanol of the desired com-
79 position χ_{MeOH} was prepared first. All reported experimental data for one
80 salt at one salt-free solvent composition χ_{MeOH} were obtained using one sin-
81 gle stock solution. The solvent (pure or mixed) was then added to a known
82 amount of salt to yield samples of about 20 ml. Uncertainties in the compo-
83 sition variables χ_{MeOH} and x_{C^+} were estimated from error propagation of the
84 uncertainty of the balance. Thereby, the uncertainty of the salt-free solvent
85 composition χ_{MeOH} is found to be better than ± 0.00002 mol mol⁻¹. The
86 uncertainty of the mole fraction of the cation x_{C^+} is found to be better than
87 ± 0.0002 mol mol⁻¹ in most cases, except for the highly concentrated CsCl

88 solutions, for which it is up to $\pm 0.0009 \text{ mol mol}^{-1}$.

89 As a guide to the accessible concentration range in case of solutions of NaCl,
90 the solubility data of Pinho and Macedo [14] were used. They proved to be
91 reliable during the present experiments. In case of CsCl, the solubility is only
92 known for the pure solvents water [27] and methanol [28], but not for mixed
93 solvents. As an estimate, we assumed the same qualitative dependence of
94 the solubility on the composition of the solvent mixture as for NaCl.

95 The densities of the samples were measured with a vibrating tube densimeter
96 (DMA 4500 M, Anton Paar), which was calibrated with air and deionized
97 water. Based on the repetition of several experiments and the resolution
98 provided by the densimeter, the uncertainty of the reported densities is es-
99 timated to be better than $\pm 0.0001 \text{ g cm}^{-3}$. The temperature was measured
100 with the densimeter's built-in thermometer, for which the supplier claims an
101 uncertainty of $\pm 0.1 \text{ K}$.

102 To obtain the partial molar volume of the salt at infinite dilution v_{salt}^{∞} in the
103 solvent from the measured densities, a simple empirical correlation is devel-
104 oped in the present work, cf. Sections 4.1 and 4.2. The uncertainty in the
105 numbers for v_{salt}^{∞} obtained from that correlation was estimated by randomly
106 disturbing the measured densities with their uncertainty and using these val-

107 ues for the fits. Thereby, the uncertainty in v_{salt}^{∞} is estimated to be better
108 than $\pm 0.5 \text{ cm}^3 \text{ mol}^{-1}$ and thus of similar magnitude as the uncertainties
109 reported by Takenaka et al. [1].

110 **3. Molecular Simulation**

111 In the present work, rigid, non-polarizable molecular models for the solvents
112 and ions are employed. The water model SPC/E is taken from the literature
113 [29], the ion models and the methanol model are taken from previous work
114 of our group. For methanol, we employ the molecular model of Schnabel et
115 al. [30], which was optimized with respect to the vapor-liquid equilibrium of
116 pure methanol. For the ions, molecular models of the Lennard-Jones (LJ)
117 + point charge type from the ion model set of Reiser et al. [31] are used.
118 The models of that set were trained together with the SPC/E water model
119 mainly using density data of dilute aqueous solutions of all alkali halide salts
120 [32]. Thus, the models used here for NaCl and CsCl were not optimized
121 for describing the individual salts, but taken from an ion model set which
122 includes all alkali and halide ions. Therefore, the Cl^- model is the same for
123 both salts studied here.

124 Molecular dynamics (MD) simulations of methanol-water mixtures, employ-

125 ing the Schnabel et al. [30] model for methanol and several popular water
 126 models (including SPC/E) were previously conducted by Guevara-Carrión
 127 et al. [33, 34] and Pařez et al. [35], but with a focus on transport proper-
 128 ties. In another previous study [36], the predictions obtained from combining
 129 the ion models of the set of Reiser et al. [31] with the methanol model of
 130 Schnabel et al. [30] using the Lorentz-Berthelot combining rules were already
 131 studied. Good agreement with experimental data for solutions of all alkali
 132 halide salts in methanol was found even though the ion models had only
 133 been trained with data on aqueous solutions. Altogether, these results from
 134 previous studies suggest that the employed models are suited as a starting
 135 point for modeling ternary solutions.
 136 All models employ LJ sites and point or partial charges, so that the potential
 137 writes [37]

$$\begin{aligned}
 U &= U_{\text{LJ}} + U_{\text{C}} \\
 &= \sum_{i=1}^{N-1} \sum_{j=i+1}^N \left\{ \sum_{a=1}^{n_i^{\text{LJ}}} \sum_{b=1}^{n_j^{\text{LJ}}} 4\epsilon_{ijab} \left[\left(\frac{\sigma_{ijab}}{r_{ijab}} \right)^{12} - \left(\frac{\sigma_{ijab}}{r_{ijab}} \right)^6 \right] \right. \\
 &\quad \left. + \sum_{c=1}^{n_i^e} \sum_{d=1}^{n_j^e} \frac{1}{4\pi\epsilon_0} \frac{q_{ic}q_{jd}}{r_{ijcd}} \right\}, \tag{3}
 \end{aligned}$$

138 where the indices a, b, c , and d refer to model interaction sites and i and j
 139 refer to molecules (including ions), ϵ_0 is the vacuum permittivity, ϵ_{ijab} and
 140 σ_{ijab} are the Lennard-Jones energy and size parameters, r_{ijab} and r_{ijcd} are
 141 site-site distances, and q_{ic} and q_{jd} are the magnitudes of the point charges.
 142 The interaction between unlike LJ sites is described by the modified Lorentz-
 143 Berthelot combining rules [38, 39]

$$\sigma_{ijab} = \eta_{ij} \frac{\sigma_{iiaa} + \sigma_{jjbb}}{2}, \quad (4)$$

$$\epsilon_{ijab} = \sqrt{\epsilon_{iiaa}\epsilon_{jjbb}}. \quad (5)$$

144 A binary interaction parameter η_{ij} different from 1 is only employed in the
 145 unlike interaction water-methanol and adjusted to experimental data of the
 146 molar excess volume at 298.15 K, cf. Section 4.1. This correction is in-
 147 troduced into the Lorentz rule here as the unlike size parameter obviously
 148 shows a much larger influence on densities than the unlike energy parameter,
 149 which is in line with a general theoretical analysis by Fischer et al. [40]. The
 150 adjustment was carried out to ensure a good description of the volumetric
 151 properties of the salt-free solvent mixture. It has only a minor impact on the

152 results obtained for the properties of the electrolytes in the mixed solvent on
153 which we focus here.

154 Densities were obtained by standard MD simulations in the NpT ensemble
155 for the temperatures 288.15, 298.15 and 318.15 K. Simulation details are
156 given in Appendix A. For each studied salt-free solvent composition, the
157 simulations were carried up to the experimental solubility limit of the salt.
158 To obtain the partial molar volume of the salt at infinite dilution in the sol-
159 vent from the densities obtained in these simulations, the same correlation
160 as for the measured densities was employed, cf. Sections 4.1 and 4.2.

161 **4. Results and Discussion**

162 The solution densities obtained from the experiments and the molecular sim-
163 ulations are reported in Tables 1 - 4. The densities of the studied salt-free
164 solvents are reported together with the data for NaCl. To improve the read-
165 ability of the plots, in most figures we only display the experimental results
166 for 288.15, 298.15 and 318.15 K, and omit those for 293.15 and 308.15 K.

Table 1: Experimental data for the density of solutions of NaCl in mixtures of water and methanol at 1 bar. The uncertainties are: $u(\chi_{\text{MeOH}}) = \pm 0.00002 \text{ mol mol}^{-1}$, $u(\rho) = \pm 0.0001 \text{ g cm}^{-3}$, $u(T) = \pm 0.1 \text{ K}$. For x_{Na^+} , the uncertainty of the last digit is given in parentheses, or omitted where it is below $\pm 0.0001 \text{ mol mol}^{-1}$.

		$\rho / \text{g cm}^{-3}$				
$\chi_{\text{MeOH}} / \text{mol mol}^{-1}$	$x_{\text{Na}^+} / \text{mol mol}^{-1}$	T / K				
		288.15	293.15	298.15	308.15	318.15
0	0	0.9991	0.9982	0.9970	0.9940	0.9902
	0.0099(1)	1.0222	1.0210	1.0196	1.0162	1.0121
	0.0196(2)	1.0447	1.0432	1.0416	1.0378	1.0335
	0.0291(2)	1.0658	1.0641	1.0623	1.0582	1.0536
	0.0385(3)	1.0867	1.0848	1.0828	1.0785	1.0737
	0.0476(3)	1.1074	1.1054	1.1033	1.0987	1.0938
	0.0566(4)	1.1272	1.1250	1.1227	1.1180	1.1129
	0.0654(4)	1.1471	1.1448	1.1424	1.1374	1.1322
0.25	0	0.9421	0.9393	0.9364	0.9301	0.9234
	0.0040	0.9503	0.9475	0.9445	0.9382	0.9315
	0.0098(1)	0.9603	0.9574	0.9544	0.9480	0.9413
	0.0229(2)	0.9845	0.9815	0.9784	0.9719	0.9651
	0.0336(2)	1.0043	1.0012	0.998	0.9914	0.9845
0.5	0	0.8892	0.8854	0.8815	0.8736	0.8653
	0.0020	0.8921	0.8883	0.8844	0.8765	0.8682
	0.0049	0.8968	0.8930	0.8891	0.8811	0.8729
	0.0079(1)	0.9014	0.8975	0.8937	0.8857	0.8775
	0.0117(1)	0.9074	0.9036	0.8997	0.8917	0.8835
	0.0146(1)	0.9125	0.9087	0.9048	0.8968	0.8886
	0.0192(1)	0.9189	0.9150	0.9111	0.9031	0.8949

Table 1 continued.

		$\rho / \text{g cm}^{-3}$				
$\chi_{\text{MeOH}} / \text{mol mol}^{-1}$	$x_{\text{Na}^+} / \text{mol mol}^{-1}$	T / K				
		288.15	293.15	298.15	308.15	318.15
0.75	0	0.8398	0.8355	0.8312	0.8224	0.8134
	0.0013	0.8418	0.8375	0.8332	0.8245	0.8155
	0.0027	0.8437	0.8395	0.8352	0.8264	0.8175
	0.0053	0.8471	0.8428	0.8385	0.8298	0.8209
	0.0079	0.8509	0.8467	0.8424	0.8337	0.8248
	0.0104(1)	0.8558	0.8516	0.8473	0.8387	0.8298
	0.0130(1)	0.8584	0.8541	0.8498	0.8412	0.8323
1	0	0.7959	0.7913	0.7866	0.7771	0.7675
	0.0010	0.7974	0.7927	0.7880	0.7786	0.7690
	0.0020	0.7989	0.7943	0.7896	0.7802	0.7707
	0.0030	0.8001	0.7955	0.7908	0.7814	0.7719
	0.0040	0.8014	0.7968	0.7921	0.7827	0.7733
	0.0050	0.8029	0.7983	0.7937	0.7843	0.7748
	0.0060	0.8041	0.7995	0.7949	0.7855	0.7761
	0.0070	0.8057	0.8011	0.7964	0.7871	0.7777

Table 2: Experimental data for the density of solutions of CsCl in mixtures of water and methanol at 1 bar. The uncertainties are: $u(\chi_{\text{MeOH}}) = \pm 0.00002 \text{ mol mol}^{-1}$, $u(\rho) = \pm 0.0001 \text{ g cm}^{-3}$, $u(T) = \pm 0.1 \text{ K}$. For x_{Cs^+} , the uncertainty of the last digit is given in parentheses, or omitted where it is below $\pm 0.0001 \text{ mol mol}^{-1}$.

		$\rho / \text{g cm}^{-3}$				
$\chi_{\text{MeOH}} / \text{mol mol}^{-1}$	$x_{\text{Cs}^+} / \text{mol mol}^{-1}$	T / K				
		288.15	293.15	298.15	308.15	318.15
0	0.0099(1)	1.0697	1.0684	1.0670	1.0635	1.0592
	0.0196(2)	1.1373	1.1358	1.1341	1.1301	1.1255
	0.0292(2)	1.2033	1.2016	1.1996	1.1952	1.1902
	0.0385(3)	1.2662	1.2642	1.2620	1.2573	1.252
	0.0536(7)	1.3670	1.3646	1.3621	1.3567	1.3509
	0.0690(9)	1.4669	1.4641	1.4613	1.4553	1.4486
0.25	0.0192(3)	1.0661	1.0650	1.0623	1.0564	1.0501
	0.0370(5)	1.1664	1.1635	1.1604	1.1540	1.1471
	0.0536(7)	1.2534	1.2501	1.2466	1.2395	1.2311
0.5	0.0074	0.9248	0.9208	0.9169	0.9088	0.9004
	0.0146(1)	0.9597	0.9557	0.9517	0.9435	0.9350
	0.0215(1)	0.9916	0.9876	0.9835	0.9751	0.9665
	0.0283(2)	1.0241	1.0200	1.0158	1.0073	0.9986
	0.0349(2)	1.0564	1.0522	1.0480	1.0394	1.0305
0.75	0.0025	0.8511	0.8468	0.8424	0.8336	0.8246
	0.0050	0.8611	0.8568	0.8524	0.8436	0.8346
	0.0074	0.8712	0.8669	0.8625	0.8536	0.8445
	0.0098(1)	0.8808	0.8764	0.8720	0.8631	0.8540
	0.0122(1)	0.8901	0.8857	0.8813	0.8723	0.8632
1	0.0012	0.8008	0.7961	0.7913	0.7818	0.7723
	0.0025	0.8055	0.8008	0.7960	0.7865	0.7769
	0.0037	0.8099	0.8052	0.8005	0.7910	0.7814
	0.0049	0.8145	0.8098	0.8051	0.7955	0.7859

Table 3: Molecular simulation data for the density of solutions of NaCl in mixtures of water and methanol at 1 bar. For the density, the uncertainty of the last digit is given in parentheses.

		$\rho / \text{g cm}^{-3}$		
$\chi_{\text{MeOH}} / \text{mol mol}^{-1}$	$x_{\text{Na}^+} / \text{mol mol}^{-1}$	T / K		
		288.15	298.15	318.15
0	0	1.0042(2)	0.9996(2)	0.9879(2)
	0.02	1.0407(2)	1.0357(2)	1.0232(2)
	0.04	1.0762(2)	1.0708(2)	1.0577(2)
	0.06	1.1107(2)	1.1045(2)	1.0907(2)
0.25	0	0.9493(2)	0.9407(2)	0.9237(2)
	0.01	0.9646(2)	0.9559(2)	0.9385(2)
	0.02	0.9789(2)	0.9710(2)	0.9531(2)
	0.03	0.9945(2)	0.9867(2)	0.9679(2)
0.5	0	0.8931(2)	0.8838(2)	0.8644(2)
	0.005	0.8995(2)	0.8904(2)	0.8710(2)
	0.01	0.9061(2)	0.8966(2)	0.8778(2)
	0.015	0.9127(2)	0.9036(2)	0.8838(2)
0.75	0	0.8427(2)	0.8337(2)	0.8141(2)
	0.004	0.8484(2)	0.8391(2)	0.8196(2)
	0.008	0.8533(2)	0.8436(2)	0.8246(2)
	0.12	0.8584(2)	0.8489(2)	0.8299(2)
1	0	0.7974(2)	0.7880(1)	0.7687(1)
	0.002	0.8002(2)	0.7910(2)	0.7717(1)
	0.004	0.8029(1)	0.7938(1)	0.7743(1)
	0.006	0.8056(2)	0.7963(2)	0.7770(1)

Table 4: Molecular simulation data for the density of solutions of CsCl in mixtures of water and methanol at 1 bar. For the density, the uncertainty of the last digit is given in parentheses.

		$\rho / \text{g cm}^{-3}$		
$\chi_{\text{MeOH}} / \text{mol mol}^{-1}$	$x_{\text{Cs}^+} / \text{mol mol}^{-1}$	T / K		
		288.15	298.15	318.15
0	0.02	1.1262(2)	1.1212(2)	1.1086(2)
	0.04	1.2437(2)	1.2379(2)	1.2245(2)
	0.06	1.3549(2)	1.3487(2)	1.3353(2)
0.25	0.02	1.0493(2)	1.0404(2)	1.0226(2)
	0.04	1.1470(2)	1.1378(3)	1.1194(2)
	0.06	1.2420(2)	1.2331(3)	1.2143(2)
0.5	0.01	0.9358(2)	0.9256(2)	0.9061(2)
	0.02	0.9772(2)	0.9678(2)	0.9479(2)
	0.03	1.0182(3)	1.0100(2)	0.9900(2)
0.75	0.004	0.8576(2)	0.8486(2)	0.8287(2)
	0.008	0.8720(2)	0.8627(2)	0.8432(2)
	0.012	0.8869(2)	0.8775(2)	0.8578(2)
1	0.002	0.8041(2)	0.7943(1)	0.7751(1)
	0.004	0.8105(1)	0.8012(1)	0.7818(1)
	0.006	0.8175(1)	0.8079(1)	0.7886(1)

167 *4.1. Salt-free Solvent Mixture*

168 Studying a ternary electrolyte solution first requires an adequate description
169 of the salt-free solvent mixture. The volumetric behavior of the salt-free
170 solvent mixture water-methanol is discussed here based on the molar excess
171 volume v^E

$$v^E(T, \chi_{\text{MeOH}}) = v(T, \chi_{\text{MeOH}}) - \chi_{\text{MeOH}} v_{\text{MeOH}}^{\text{pure}}(T) - (1 - \chi_{\text{MeOH}}) v_{\text{W}}^{\text{pure}}(T), \quad (6)$$

172 where v is the molar volume of the mixture, v_i^{pure} are the pure component
173 molar volumes, and the methanol mole fraction is χ_{MeOH} in our notation.
174 Results for the molar excess volume of the mixture water + methanol at
175 298.15 K are shown in Fig. 1.

176 The experimental results from the present work are compared to a correlation
177 of experimental data of Coquelet et al. [41]. Excellent agreement is observed.
178 The deviations are below about $\pm 0.01 \text{ cm}^3 \text{ mol}^{-1}$. Furthermore, molecular
179 simulation results from the present work are shown. They were obtained
180 using $\eta_{\text{W-MeOH}} = 0.993$. That parameter was fit to reproduce the minimum
181 of v^E , which is found for equimolar composition both experimentally and




Figure 1: Molar excess volume of mixtures of water and methanol at 298.15 K. (\square) experimental data, this work, (\circ) molecular simulation data, this work, (—) correlation of experimental data by Coquelet et al. [41], (---) correlation of the present molecular simulation data.

182 in the simulations. The deviation of $\eta_{\text{W-MeOH}}$ from 1 is only small. Using
 183 $\eta_{\text{W-MeOH}} = 1$, the minimum of v^{E} is about 15% above the experimental
 184 result. Using the adjusted value of $\eta_{\text{W-MeOH}}$, the dependence of v^{E} on χ_{MeOH}
 185 is predicted well by the simulation.

186 The density of the salt-free solvent mixture is the basis for the correlation
 187 used here for describing the density of the ternary electrolyte solutions. As
 188 Coquelet et al. [41], we employ a Redlich-Kister type correlation, which in
 189 our notation writes as

$$v^E(T, \chi_{\text{MeOH}}) = (1 - \chi_{\text{MeOH}})\chi_{\text{MeOH}} \sum_k A_k(T)(1 - 2\chi_{\text{MeOH}})^k, \quad k = 0, 1, \dots, \quad (7)$$

190 where the A_k are fit parameters. We also use Eq. (7) here for correlating
 191 the molecular simulation data. The Redlich-Kister coefficients A_k for de-
 192 scribing the molar excess volume of mixtures of water and methanol for all
 193 temperatures studied here are presented in Appendix B, cf. Table B.1. Two
 194 parameter sets are reported. The first one describes the experimental data.
 195 It is adopted from Coquelet et al. [41], but also describes the present exper-
 196 imental results within their uncertainty. The second one is obtained from
 197 a fit to the present simulation data for the system water + methanol. The
 198 densities of pure water and pure methanol, which are needed in addition to
 199 the Redlich-Kister fit to calculate the density of the mixture, are taken from
 200 Tables 1 and 3.

201 *4.2. Electrolyte Solutions*

202 Fig. 2 shows the measured and simulated densities of solutions of NaCl and
 203 of CsCl in mixtures of water and methanol at 298.15 K.




Figure 2: Densities of solutions of NaCl (left column) and CsCl (right column) in mixtures of water and methanol over the cation mole fraction at 298.15 K. Results from the present work: (\square) experiment, (\circ) molecular simulation, (—) correlation of experimental data, (---) correlation of molecular simulation data.

204 For both salts in pure water ($\chi_{\text{MeOH}} = 0$), the increase in the density upon
205 addition of the salt is slightly underpredicted by the models. The salt models
206 used here are not individual models but taken from an ion model set for
207 all alkali halides, so that compromises regarding the performance for some
208 salts had to be made [32]. However, the higher the methanol content in
209 the mixture, the better the agreement between experiment and simulation.
210 This is a remarkable finding since in the development of the ion models, only
211 data on aqueous solutions were used. The influence of adding methanol is
212 predicted surprisingly well by the models.

213 In previous work of our group [36, 42], it was found that both for aqueous
214 and methanolic solutions of alkali halide salts, the density of the solution is
215 an almost perfectly linear function of the ion mole fraction. That finding was
216 based on results for $x_{\text{C}^+} \leq 0.05 \text{ mol mol}^{-1}$. The present results, both from
217 experiment and from molecular simulation, show that the linearity holds also
218 up to the solubility limit for the studied salts. Additionally, the linear relation
219 is also found to hold for mixed solvents. Thus, for the ternary electrolyte
220 solutions, we employ the correlation

$$\rho(T, \chi_{\text{MeOH}}, x_{\text{C}^+}) = \rho_{\text{solv}}(T, \chi_{\text{MeOH}}) + b_{\text{salt}}(T, \chi_{\text{MeOH}}) x_{\text{C}^+} \quad (8)$$

$$\text{with } b_{\text{salt}} = \left(\frac{\partial \rho}{\partial x_{\text{C}^+}} \right)_{T, \chi_{\text{MeOH}}}, \quad (9)$$

221 where ρ_{solv} is the density of the salt-free solvent mixture, which is obtained
 222 as described in the previous section. The slope of the density b_{salt} was fit to
 223 the data for each individual solvent composition.
 224 In Fig. 3, the slope of the density b_{salt} determined from experimental and
 225 molecular simulation data is shown for 288.15, 298.15 and 318.15 K.




Figure 3: Slope of the density b_{salt} , cf. Eq. (9), of NaCl solutions (left) and CsCl solutions (right) over the composition of the salt-free solvent mixture water + methanol. Colors indicate the isotherms 288.15 K (blue), 298.15 K (gray) and 318.15 K (red). As the temperature influence is small, the results overlap. Results from the present work: (\square) experiment, (\circ) molecular simulation, (—) correlation of experimental data, (---) correlation of molecular simulation data.

226 The experimental data of the NaCl solutions reveal a surprising behavior:
227 For aqueous solutions, the slope of the density decreases with increasing
228 temperature. This trend is reversed for methanolic solutions. Furthermore,
229 at approximately equimolar composition of the salt-free solvent mixture, the
230 slope of the density is independent of the temperature. This behavior is also
231 predicted by the molecular simulations. The experimental results show that
232 b_{NaCl} does not decline linearly with increasing χ_{MeOH} . The trend is predicted
233 well by the molecular simulations. In the simulations, a shallow minimum is
234 found for all temperatures, which is present in the experimental data only
235 for the higher temperatures.

236 The experimental data of the CsCl solutions show a more steady decline of the
237 slope of the density when adding methanol to the solution. Comparing NaCl
238 and CsCl in pure methanol, the temperature dependence is reversed: b_{NaCl}
239 decreases with increasing temperature, while b_{CsCl} increases with increasing
240 temperature. Consequently, for CsCl solutions the isotherms of the slope of
241 the density do not intersect. In general, the temperature dependence of b_{CsCl}
242 is weaker than the temperature dependence of b_{NaCl} . All this is predicted
243 well by the molecular models.

244 To establish a correlation, b_{salt} is described by a polynomial of second degree:

$$b_{salt}(T, \chi_{\text{MeOH}}) = b_{2,salt}(T)\chi_{\text{MeOH}}^2 + b_{1,salt}(T)\chi_{\text{MeOH}} + b_{0,salt}(T) \quad (10)$$

245 The fit parameters $b_{k,salt}$ were determined individually for each isotherm for
 246 both the experimental and the simulation data and are given in Appendix
 247 B, cf. Table B.2.

248 Together with the pure component molar volumes of water and methanol,
 249 Eqs. (7), (8) and (10) fully determine the density in the ternary system
 250 water-methanol-salt. From that correlation, it is possible to deduce any
 251 volumetric property of interest. In the following, we focus on the partial
 252 molar volume of the salt at infinite dilution v_{salt}^∞ in the mixed solvent with
 253 composition χ_{MeOH} . It can be shown that, from the correlation developed
 254 here, this quantity can be obtained as

$$v_{salt}^\infty(T, \chi_{\text{MeOH}}) = \frac{-b_{salt}(T, \chi_{\text{MeOH}}) M_{solv} + M_{salt} \rho_{solv}(T, \chi_{\text{MeOH}})}{\rho_{solv}(T, \chi_{\text{MeOH}})^2} \quad (11)$$

$$M_{solv} = \chi_{\text{MeOH}} M_{\text{MeOH}} + (1 - \chi_{\text{MeOH}}) M_{\text{W}}, \quad (12)$$

255 where M_i is the molar mass of component i .

256 The results for the partial molar volume of both salts at infinite dilution as

257 determined from experiments and predicted by the simulations are shown in

258 Fig. 4.




Figure 4: Partial molar volume of the salt at infinite dilution v_{salt}^{∞} , cf. Eq. (11), of NaCl solutions (left) and CsCl solutions (right) over the composition of the salt-free solvent mixture water + methanol. Colors indicate the isotherms 288.15 K (blue), 298.15 K (gray) and 318.15 K (red). Results from the present work: (—) correlation of experimental data, (---) correlation of molecular simulation data.

259 Both for NaCl and CsCl, the partial molar volume of the salt is distinctly
260 lower in methanol than in water. This indicates that the ions have a strong
261 ordering effect especially on methanol. For NaCl solutions, v_{NaCl}^{∞} depends
262 only weakly on χ_{MeOH} for χ_{MeOH} below about 0.5 mol mol⁻¹, while for larger
263 χ_{MeOH} an important decrease of v_{NaCl}^{∞} is observed. For CsCl solutions, the
264 decrease of v_{CsCl}^{∞} is more steady.

265 The temperature dependence of v_{salt}^{∞} is weak both for NaCl and CsCl. While
266 v_{NaCl}^{∞} increases slightly with increasing temperature in water, it decreases

267 with increasing temperature in methanol. The same is true for v_{CsCl}^{∞} . For
268 both salts, there is a solvent composition for which the temperature depen-
269 dence vanishes. For high temperatures, the partial molar volume of NaCl
270 in methanol is negative. This is remarkable since despite addition of salt,
271 the volume of the solution decreases. In contrast, the partial molar volume
272 of CsCl in methanol is positive. Comparing both salts and considering that
273 they comprise the same anion shows that especially Na^+ ions have a strong
274 influence on the structure of methanol.

275 The experimental findings discussed above are correctly predicted by the
276 molecular simulations. However, there are some quantitative differences
277 which are of the order of $10 \text{ cm}^3 \text{ mol}^{-1}$ for both salts. Fig. 4 clearly shows
278 that they are induced by the models of the systems salt + pure solvent, while
279 the effects of the mixed solvents are well predicted. Taking into account that
280 the salt models were taken from an ion model set and not adjusted individ-
281 ually and the fact that adjustments of the ion models were only made using
282 experimental data for the solvent water, the predictions are of remarkable
283 quality.

284 To point out the different behavior of the salts in the two pure solvents water
285 and methanol more clearly, the temperature dependence of v_{salt}^{∞} in both pure

286 solvents is shown in Fig. 5.




Figure 5: Partial molar volume of the salt at infinite dilution v_{salt}^{∞} , cf. Eq. (11), of NaCl solutions (left) and CsCl solutions (right) over the temperature. The two sets of data in each plot correspond to pure water as the solvent ($\chi_{\text{MeOH}} = 0$, top) and pure methanol as the solvent ($\chi_{\text{MeOH}} = 1$, bottom). Results from the present work: (\square) experiment, (\circ) molecular simulation. Dotted lines are guides to the eye.

287 As already known from Fig. 4, for both salts v_{salt}^{∞} increases with increasing
 288 temperature in an aqueous solution, while it decreases with increasing tem-
 289 perature in a methanolic solution. This feature and also the quantitative
 290 incline / decline is predicted well by the molecular simulations. It is instruc-
 291 tive to interpret this behavior in the light of Kirkwood-Buff theory [43–45].
 292 The present results indicate that for both salts in water, the total correlation
 293 of the ions with the solvent molecules is lowered with increasing temperature.
 294 In contrast, for both salts in methanol the total correlation of the ions with
 295 the solvent molecules is enhanced with increasing temperature. A rigorous

296 assessment of this relation might be carried out using additional simulations.
297 However, that study is beyond the scope of the present work as large system
298 sizes are needed to avoid finite size effects [46] and Kirkwood-Buff integrals
299 usually show poor convergence [47], which is especially cumbersome for the
300 infinite dilution case studied here.

301 *4.3. Comparison of Experimental Data from the Literature and the Present* 302 *Work*

303 For both systems studied in the present work, density data have been re-
304 ported before.

305 There are four sets of data for NaCl in solutions of water and methanol:
306 Takenaka et al. [1] studied the system in great detail, covering the same con-
307 centration and temperature ranges as the present work. Khimenko [4] and
308 Werblan [5] studied only the isotherms 293.15 K and 298.15 K, respectively,
309 and their works also cover only parts of the concentration range. For these
310 two sets of data, the numbers were taken here from the Dortmund Data Bank
311 [48] because the original sources could not be retrieved. The fourth data set
312 on NaCl solutions is that of Guetachew et al. [6], who report densities at
313 298.15 K. Unfortunately, it is difficult to compare their data to any of the
314 other sets because their measurements were not carried out at constant com-

315 positions of the salt-free solvent mixture. That data set is therefore omitted
316 in the following discussion.

317 For solutions of CsCl in mixtures of water and methanol, densities have so
318 far only been reported by Werblan [5]. Also that data set was taken from
319 the Dortmund Data Bank because the original source could not be retrieved.

320 In Fig. 6, the experimental data from the literature are compared to the cor-
321 relation of experimental data developed in the present work. Since different
322 salt-free solvent compositions were studied in the literature, this comparison
323 is carried out based on the slope of the density $b_{salt}(\chi_{\text{MeOH}})$ at 298.15 K. To
324 this end, the literature data were correlated using the same linear relation-
325 ship as for the data of the present work, cf. Eq. (8). The linear trend was
326 confirmed from inspection of the data of Takenaka et al. [1] and of Khimenko
327 [4], while the data of Werblan [5] showed considerable scatter.

328 For the pure solvents water and methanol, the results from experiments from
329 previous work of our group [36, 42] are also included in Fig. 6. They are
330 in very good agreement with the correlation developed in the present work.

331 For solutions of NaCl in mixtures of water and methanol, the agreement
332 between the data of Takenaka et al. [1] and the correlation to experimental
333 data obtained in the present work is excellent. The data of Khimenko [4],




Figure 6: Slope of the density b_{salt} , cf. Eq. (9), of NaCl solutions (left) and CsCl solutions (right) over the composition of the salt-free solvent mixture water + methanol. All displayed data are at 298.15 K except for the results of Khimenko [4], which are at 293.15 K. The lines represent the correlation of experimental data from the present work. Symbols denote experimental data from the literature: (\diamond) Takenaka et al. [1], (\times) Khimenko [4], (\star) Werblan [5], (\triangle) Reiser et al. [36, 42]. For solutions of NaCl and CsCl in pure water, the results of Reiser et al. [42] were interpolated between 293.15 and 303.15 K.

334 which were reported only for high water concentrations, deviate considerably
 335 from both our data and those of Takenaka et al. [1]. The same holds for most
 336 of the data of Werblan [5], which scatter very strongly.

337 5. Conclusions

338 Densities of solutions of NaCl and CsCl in mixtures of water and methanol
 339 were studied by experiment and molecular simulation. For the NaCl solu-
 340 tions, the experimental data from the present work are found to be in very
 341 good agreement to those reported by Takenaka et al. [1]. For the CsCl solu-

342 tions, up to now only the unreliable data of Werblan [5] were available.

343 The employed molecular models for the ions were taken from a set of models
344 for all alkali and halide ions, for which the parameters were obtained by a
345 fit to density data of dilute aqueous solutions only. The ion models are com-
346 bined with established solvent models using the Lorentz-Berthelot combining
347 rules. The models show excellent predictions for the dependence of the elec-
348 trolyte solution density on the composition of the salt-free solvent mixture.

349 Also the temperature dependence is predicted correctly. Some quantitative
350 differences are found, which stem, however, from deviations in the models
351 of the electrolytes in the pure solvents, for which no individual adjustments
352 were carried out. Altogether, taking into account that the employed molec-
353 ular models are very simple, the results are very encouraging.

354 Both salts are found to behave differently in the two pure solvents water and
355 methanol. The present results indicate that the effect of ions on methanol
356 is even more pronounced than their effect on water. This is especially true
357 for the Na^+ ion. The good agreement between simulations and experiments
358 shows that the employed models give a realistic picture of the solution be-
359 havior. Therefore, subsequent molecular simulations should be carried out in
360 future work, as they can provide additional insight into the solution behavior

361 on the molecular level.

362 **Acknowledgments**

363 The authors gratefully acknowledge financial support by the Reinhart Kosel-
364 leck Program (HA1993/15-1) of the German Research Foundation (DFG).

365 The present work was conducted under the auspices of the Boltzmann-Zuse
366 Society of Computational Molecular Engineering (BZS) and the simulations
367 were carried out on the Regional University Computing Center Kaisers-
368 lautern (RHRK) under the grant TUKL-TLMV, the High Performance Com-
369 puting Center Stuttgart (HLRS) under the grant MMHBF as well as the Leib-
370 nitz Supercomputing Centre (LRZ) under the grant SPARLAMPE (pr48te).

371 **Funding**

372 This work was supported by the Reinhart Koselleck Program (HA1993/15-1)
373 of the German Research Foundation (DFG).

374 **Appendix A. Simulation Details**

375 In the present work, MD simulations were carried out with the molecular
376 simulation program *ms2* [49]. Equilibration and production took 100,000
377 and 2,000,000 time steps of length 1.2 fs, respectively. The equations of

378 motion were integrated with a gear predictor-corrector scheme of fifth or-
379 der. The simulations were run with a total of 1000 particles in the NpT
380 ensemble, employing the velocity scaling thermostat and Andersen’s baro-
381 stat. All interactions were evaluated explicitly up to the cutoff radius of 15
382 Å. The usual LJ long-range corrections to the virial and the energy were
383 included. Electrostatic long-range interactions were calculated using Ewald
384 summation, where the real and reciprocal space cutoff were 15 Å. Statistical
385 simulation uncertainties were estimated with the block average method by
386 Flyvbjerg and Petersen [50].

387 **Appendix B. Correlation Parameters**

388 Tables B.1 and B.2 contain the parameters of the empirical density correla-
389 tion developed in the present work.

Table B.1: Parameters of the Redlich-Kister fits for the molar excess volume of mixtures of water and methanol, cf. Eq. (7).

T / K	Experiment ^a						Simulation ^b		
	A_0	A_1	A_2	A_3	A_4	A_5	A_0	A_1	A_2
288.15	-4.024	-0.319	0.113	0.382	0.621	0.291	-4.147	0.359	-0.387
293.15	-4.047	-0.314	0.125	0.462	0.475	-0.073	—	—	—
298.15	-4.069	-0.299	0.121	0.492	0.388	-0.283	-4.094	0.152	-0.492
308.15	-4.111	-0.219	0.082	0.378	0.328	-0.254	—	—	—
318.15	-4.146	-0.090	0.001	0.105	0.437	0.235	-4.020	0.139	-0.697

^aTaken from Coquelet et al. [41]

^bNo simulations were carried out for 293.15 K and 308.15 K

Table B.2: Parameters of the polynomial fits to the slope of the density, cf. Eq. (10).

NaCl						
	Experiment			Simulation ^a		
T / K	$b_{2,\text{NaCl}}$	$b_{1,\text{NaCl}}$	$b_{0,\text{NaCl}}$	$b_{2,\text{NaCl}}$	$b_{1,\text{NaCl}}$	$b_{0,\text{NaCl}}$
288.15	1.0506	-1.9072	2.2658	0.9993	-1.3956	1.7788
293.15	1.0167	-1.8504	2.2444	—	—	—
298.15	0.9843	-1.7991	2.2255	0.9330	-1.3336	1.7662
308.15	0.9880	-1.7470	2.1975	—	—	—
318.15	0.9829	-1.6953	2.1778	0.9654	-1.2788	1.7172
CsCl						
	Experiment			Simulation ^a		
T / K	$b_{2,\text{CsCl}}$	$b_{1,\text{CsCl}}$	$b_{0,\text{CsCl}}$	$b_{2,\text{CsCl}}$	$b_{1,\text{CsCl}}$	$b_{0,\text{CsCl}}$
288.15	1.9178	-5.0312	6.8499	1.6940	-4.1679	5.8345
293.15	1.8580	-4.9636	6.8225	—	—	—
298.15	1.8465	-4.9312	6.8002	1.5320	-4.0245	5.8094
308.15	1.7835	-4.8498	6.7596	—	—	—
318.15	1.7780	-4.8008	6.7170	1.5604	-4.0167	5.7797

^aNo simulations were carried out for 293.15 K and 308.15 K

390 **Nomenclature**

391 *Abbreviations*

EOS Equation of state

LJ Lennard-Jones

MeOH Methanol

MD Molecular dynamics

W Water

392 *Symbols*

b_{salt} Slope of the density versus ion mole fraction

ϵ LJ energy parameter

ϵ_0 Vacuum permittivity

n_i Number of moles of component i

n_i^j Number of sites of type j on molecule i

N Number of molecules

η_{ij} Binary interaction parameter

M_i Molar mass of component i

q Point charge

ρ Mass density

r Distance

σ	LJ size parameter
T	Temperature
$u(z)$	Uncertainty of the property z
U	Potential
v	Molar volume
v^E	Molar excess volume
v_i	Partial molar volume of component i
v_i^{pure}	Molar volume of pure component i
x_i	Mole fraction of component i
χ_{MeOH}	Methanol mole fraction in the salt-free solvent mixture

393 *Subscripts and Superscripts*

a, b, c, d	Site index
e	Electrostatic
i, j	Component / molecule index
k	Index for fit parameters
salt	Either NaCl or CsCl
solv	Solvent
∞	At infinite dilution

394 **References**

- 395 [1] N. Takenaka, T. Takemura, M. Sakurai, Partial Molar Volumes of Uni-
396 Univalent Electrolytes in Methanol plus Water 1: Lithium Chloride,
397 Sodium Chloride, and Potassium Chloride, *J. Chem. Eng. Data* 39
398 (1994) 207–213. doi:10.1021/je00014a004.
- 399 [2] N. Takenaka, T. Takemura, M. Sakurai, Partial Molar Volumes of
400 Uni-Univalent Electrolytes in Methanol plus Water 2: Sodium Bro-
401 mide and Potassium Bromide, *J. Chem. Eng. Data* 39 (1994) 796–801.
402 doi:10.1021/je00016a036.
- 403 [3] N. Takenaka, T. Takemura, M. Sakurai, Partial Molal Volumes of
404 Uni-Univalent Electrolytes in Methanol plus Water 3. Sodium Io-
405 dide and Potassium Iodide, *J. Chem. Eng. Data* 39 (1994) 802–807.
406 doi:10.1021/je00016a037.
- 407 [4] M. Khimenko, *Russ. J. Phys. Chem.* 43 (1969) 1043–1046.
- 408 [5] L. Werblan, *Bull. Acad. Pol. Sci. Ser. Sci. Chim.* 27 (1979) 873–890.
- 409 [6] T. Guetachew, S. Ye, I. Mokbel, J. Jose, P. Xans, Study of NaCl solu-
410 tions in a mixed solvent H₂O-CH₃OH: Experimental densities and com-

- 411 parison with calculated values obtained with a modified Pitzer's model,
412 J. Solution Chem. 25 (1996) 895–903. doi:10.1007/BF00972580.
- 413 [7] E. Ivanov, V. Abrosimov, Volumetric properties of a stoichiometric mix-
414 ture of K^+ and Br^- ions in H/D isotope-substituted aqueous methanol
415 at 278.15-318.15 K: I. The $H_2O-CH_3OH-KBr$ system, Russ. J. Inorg.
416 Chem. 51 (2006) 662–669. doi:10.1134/S0036023606040279.
- 417 [8] W. Raatschen, Thermophysikalische Eigenschaften von Methanol /
418 Wasser - Lithiumbromid Lösungen, Deutscher Kälte und Klimatechnis-
419 cher Verein e.V., Stuttgart, 1985.
- 420 [9] A. Gupta, Thermodynamics of Electrolytes in Mixed Solvents - Appli-
421 cation of Pitzer's Thermodynamic Equations to Activity Coefficients of
422 1-1 Electrolytes in Methanol-Water Mixtures, J. Phys. Chem. 83 (1979)
423 2986–2990. doi:10.1021/j100486a010.
- 424 [10] B. Mock, L. Evans, C. Chen, Thermodynamic Representation of Phase
425 Equilibria of Mixed-Solvent Electrolyte Systems, AIChE J. 32 (1986)
426 1655–1664. doi:10.1002/aic.690321009.
- 427 [11] I. Kikic, M. Fermeglia, P. Rasmussen, UNIFAC Prediction of Vapor-

- 428 Liquid-Equilibria in Mixed-Solvent Salt Systems, *Chem. Eng. Sci.* 46
429 (1991) 2775–2780. doi:10.1016/0009-2509(91)85146-O.
- 430 [12] H. J. Strauch, P. T. Cummings, Gibbs Ensemble Simulation of Mixed-
431 Solvent Electrolyte Solutions, *Fluid Phase Equilib.* 86 (1993) 147–172.
432 doi:10.1016/0378-3812(93)87173-X.
- 433 [13] C. Achard, C. Dussap, J. Gros, Representation of Vapor-Liquid-
434 Equilibria in Water-Alcohol Electrolyte Mixtures with a Modified UNI-
435 FAC Group-Contribution Method, *Fluid Phase Equilib.* 98 (1994) 71–89.
436 doi:10.1016/0378-3812(94)80109-6.
- 437 [14] S. Pinho, E. Macedo, Representation of salt solubility in mixed sol-
438 vents: A comparison of thermodynamic models, *Fluid Phase Equilib.*
439 116 (1996) 209–216. doi:10.1016/0378-3812(95)02889-7.
- 440 [15] M. Iliuta, K. Thomsen, P. Rasmussen, Extended UNIQUAC model for
441 correlation and prediction of vapour-liquid-solid equilibria in aqueous
442 salt systems containing non-electrolytes. Part A. Methanol-water-salt
443 systems, *Chem. Eng. Sci.* 55 (2000) 2673–2686. doi:10.1016/S0009-
444 2509(99)00534-5.
- 445 [16] F. Deyhimi, Z. Karimzadeh, M. Abedi, Pitzer and Pitzer-Simonson-

- 446 Clegg ion-interaction modeling approaches: Ternary HCl plus methanol
447 plus water electrolyte system, *J. Mol. Liq.* 150 (2009) 62–67.
448 doi:10.1016/j.molliq.2009.09.014.
- 449 [17] C. Held, A. Prinz, V. Wallmeyer, G. Sadowski, Measuring and
450 modeling alcohol/salt systems, *Chem. Eng. Sci.* 68 (2012) 328–339.
451 doi:10.1016/j.ces.2011.09.040.
- 452 [18] J. Schreckenber, S. Dufal, A. Haslam, C. Adjiman, G. Jackson,
453 A. Galindo, Modelling of the thermodynamic and solvation proper-
454 ties of electrolyte solutions with the statistical associating fluid the-
455 ory for potentials of variable range, *Mol. Phys.* 112 (2014) 2339–2364.
456 doi:10.1080/00268976.2014.910316.
- 457 [19] B. Maribo-Mogensen, K. Thomsen, G. Kontogeorgis, An electrolyte
458 CPA equation of state for mixed solvent electrolytes, *AIChE J.* 61 (2015)
459 2933–2950. doi:10.1002/aic.14829.
- 460 [20] G. Lamoureux, B. Roux, Absolute hydration free energy scale for alkali
461 and halide ions established from simulations with a polarizable force
462 field, *J. Phys. Chem. B* 110 (2006) 3308–3322. doi:10.1021/jp056043p.

- 463 [21] P. Kiss, A. Baranyai, A new polarizable force field for alkali and halide
464 ions, *J. Chem. Phys.* 141 (2014) 114501. doi:10.1063/1.4895129.
- 465 [22] H. Jiang, Z. Mester, O. Moulτος, I. Economou, A. Panagiotopoulos,
466 Thermodynamic and Transport Properties of H₂O + NaCl from Po-
467 larizable Force Fields, *J. Chem. Theory Comput.* 11 (2015) 3802–3810.
468 doi:10.1021/acs.jctc.5b00421.
- 469 [23] F. Moučka, I. Nezbeda, W. Smith, Chemical Potentials, Activity Coef-
470 ficients, and Solubility in Aqueous NaCl Solutions: Prediction by Po-
471 larizable Force Fields, *J. Chem. Theory Comput.* 11 (2015) 1756–1764.
472 doi:10.1021/acs.jctc.5b00018.
- 473 [24] G. Orozco, O. Moulτος, H. Jiang, I. Economou, A. Panagiotopou-
474 los, Molecular simulation of thermodynamic and transport proper-
475 ties for the H₂O+NaCl system, *J. Chem. Phys.* 141 (2014) 234507.
476 doi:10.1063/1.4903928.
- 477 [25] M. Kohns, M. Schappals, M. Horsch, H. Hasse, Activities in Aqueous
478 Solutions of the Alkali Halide Salts from Molecular Simulation, *J. Chem.*
479 *Eng. Data* 61 (12) (2016) 4068–4076. doi:10.1021/acs.jced.6b00544.
- 480 [26] I. Nezbeda, F. Moucka, W. Smith, Recent progress in molec-

- 481 ular simulation of aqueous electrolytes: force fields, chemi-
482 cal potentials and solubility, *Mol. Phys.* 114 (2016) 1665–1690.
483 doi:10.1080/00268976.2016.1165296.
- 484 [27] D. Lide (Ed.), *CRC Handbook of Chemistry and Physics, Internet Ver-*
485 *sion 2005*, CRC Press, Boca Raton, 2005.
- 486 [28] V. Stenger, Solubilities of various alkali metal and alkaline earth metal
487 compounds in methanol, *J. Chem. Eng. Data* 41 (1996) 1111–1113.
488 doi:10.1021/je960124k.
- 489 [29] H. Berendsen, J. Grigera, T. Straatsma, The Missing Term in
490 Effective Pair Potentials, *J. Phys. Chem.* 91 (1987) 6269–6271.
491 doi:10.1021/j100308a038.
- 492 [30] T. Schnabel, A. Srivastava, J. Vrabec, H. Hasse, Hydrogen bonding of
493 methanol in supercritical CO₂: Comparison between H-1 NMR spec-
494 troscopic data and molecular simulation results, *J. Phys. Chem. B* 111
495 (2007) 9871–9878. doi:10.1021/jp0720338.
- 496 [31] S. Reiser, S. Deublein, J. Vrabec, H. Hasse, Molecular dispersion energy
497 parameters for alkali and halide ions in aqueous solution, *J. Chem. Phys.*
498 140 (2014) 044504. doi:10.1063/1.4858392.

- 499 [32] S. Deublein, J. Vrabec, H. Hasse, A set of molecular models for alkali
500 and halide ions in aqueous solution, *J. Chem. Phys.* 136 (2012) 084501.
501 doi:10.1063/1.3687238.
- 502 [33] G. Guevara-Carrión, J. Vrabec, H. Hasse, Prediction of self-diffusion
503 coefficient and shear viscosity of water and its binary mixtures with
504 methanol and ethanol by molecular simulation, *J. Chem. Phys.* 134
505 (2011) 074508. doi:10.1063/1.3515262.
- 506 [34] G. Guevara-Carrión, Y. Gaponenko, T. Janzen, J. Vrabec,
507 V. Shevtsova, Diffusion in Multicomponent Liquids: From Microscopic
508 to Macroscopic Scales, *J. Phys. Chem. B* 120 (2016) 12193–12210.
509 doi:10.1021/acs.jpcc.6b09810.
- 510 [35] S. Pařez, G. Guevara-Carrión, H. Hasse, J. Vrabec, Mutual diffusion
511 in the ternary mixture of water plus methanol plus ethanol and its
512 binary subsystems, *Phys. Chem. Chem. Phys.* 15 (2013) 3985–4001.
513 doi:10.1039/c3cp43785j.
- 514 [36] S. Reiser, M. Horsch, H. Hasse, Density of Methanolic Alkali Halide Salt
515 Solutions by Experiment and Molecular Simulation, *J. Chem. Eng. Data*
516 60 (2015) 1614–1628. doi:10.1021/je5009944.

- 517 [37] C. G. Gray, K. E. Gubbins, Theory of Molecular Fluids, Vol. 1: Funda-
518 mentals, Clarendon Press, Oxford, 1984.
- 519 [38] H. Lorentz, Ueber die Anwendung des Satzes vom Virial in der kinetis-
520 chen Theorie der Gase, Ann. Phys. 248 (1881) 127–136.
- 521 [39] D. Berthelot, Sur le melange des gaz, Comptes Rendus de l’Academie
522 des Sciences Paris 126 (1898) 1703–1706, 1857–1858.
- 523 [40] J. Fischer, D. Moller, A. Chialvo, J. Haile, The Influence of Un-
524 like Molecule Interaction Parameters on Liquid-Mixture Excess Prop-
525 erties, Fluid Phase Equilib. 48 (1989) 161–176. doi:10.1016/0378-
526 3812(89)80199-2.
- 527 [41] C. Coquelet, A. Valtz, D. Richon, Volumetric properties of water
528 plus monoethanolamine plus methanol mixtures at atmospheric pres-
529 sure from 283.15 to 353.15 K, J. Chem. Eng. Data 50 (2005) 412–418.
530 doi:10.1021/je049691v.
- 531 [42] S. Reiser, M. Horsch, H. Hasse, Temperature Dependence of the
532 Density of Aqueous Alkali Halide Salt Solutions by Experiment and
533 Molecular Simulation, J. Chem. Eng. Data 59 (2014) 3434–3448.
534 doi:10.1021/je5004209.

- 535 [43] J. Kirkwood, F. Buff, The Statistical Mechanical Theory of Solutions.
536 1., J. Chem. Phys. 19 (1951) 774–777. doi:10.1063/1.1748352.
- 537 [44] P. Kusalik, G. Patey, The Thermodynamic Properties of Electrolyte
538 Solutions - Some Formal Results, J. Chem. Phys. 86 (1987) 5110–5116.
539 doi:10.1063/1.452629.
- 540 [45] A. Chialvo, P. Cummings, J. Simonson, R. Mesmer, Solvation in high-
541 temperature electrolyte solutions. II. Some formal results, J. Chem.
542 Phys. 110 (1999) 1075–1086. doi:10.1063/1.478151.
- 543 [46] P. Krüger, S. Schnell, D. Bedeaux, S. Kjelstrup, T. Vlugt, J.-M. Simon,
544 Kirkwood-Buff Integrals for Finite Volumes, J. Phys. Chem. Lett. 4
545 (2013) 235–238. doi:10.1021/jz301992u.
- 546 [47] S. Schnell, P. Englebienne, J.-M. Simon, P. Krüger, S. Balaji, S. Kjel-
547 strup, D. Bedeaux, A. Bardow, T. Vlugt, How to apply the Kirkwood-
548 Buff theory to individual species in salt solutions, Chem. Phys. Lett.
549 582 (2013) 154–157. doi:10.1016/j.cplett.2013.07.043.
- 550 [48] DDBST GmbH, Dortmund Data Bank, Version 2016, DDBST GmbH,
551 Oldenburg, 2016.

- 552 [49] C. W. Glass, S. Reiser, G. Rutkai, S. Deublein, A. Köster, G. Guevara-
553 Carrion, A. Wafai, M. Horsch, M. Bernreuther, T. Windmann, H. Hasse,
554 J. Vrabec, ms2: A Molecular Simulation Tool for Thermodynamic Prop-
555 erties, New Version Release, Comput. Phys. Commun. 185 (2014) 3302–
556 3306. doi:10.1016/j.cpc.2014.07.012.
- 557 [50] H. Flyvbjerg, H. Petersen, Error-estimates on Averages of Correlated
558 Data, J. Chem. Phys. 91 (1989) 461–466. doi:10.1063/1.457480.